

CLARK COUNTY EMS SYSTEM EMERGENCY MEDICAL CARE PROTOCOLS



EFFECTIVE: August 05, 2022
(Replaces January 31, 2022 Version)

PO BOX 3902 – LAS VEGAS, NV 89127

TABLE OF CONTENTS

Foreword.....	5
Terms and Conventions.....	8
ADULT TREATMENT PROTOCOLS.....	10
General Adult Assessment	11
General Adult Trauma Assessment.....	13
Abdominal Pain/Flank Pain, Nausea & Vomiting	15
Allergic Reaction	17
Altered Mental Status/Syncope	19
Behavioral Emergencies	21
Bradycardia	23
Burns	25
Cardiac Arrest (Non-Traumatic).....	27
Chest Pain (Non-Traumatic) and Suspected Acute Coronary Syndrome.....	29
Childbirth/Labor.....	31
Cold Related Illness	33
Drowning.....	35
Epistaxis.....	37
Heat Related Illness.....	39
Hyperkalemia (Suspected)	41
Obstetrical Emergency	43
Overdose/Poisoning.....	45
Pain Management.....	47
Pulmonary Edema/CHF	49
Respiratory Distress	51
Seizure.....	53
Sepsis	55
Shock.....	57
Smoke Inhalation	59
STEMI (Suspected)	61
Stroke (CVA)	63
Tachycardia/Stable.....	65
Tachycardia/Unstable	67
Ventilation Management	69
PEDIATRIC TREATMENT PROTOCOLS.....	71
General Pediatric Assessment.....	72
General Pediatric Trauma Assessment	74
Abdominal/Flank Pain, Nausea & Vomiting	76
Allergic Reaction	78
Altered Mental Status	80
Behavioral Emergencies	82
Bradycardia	84
Burns	86

PEDIATRIC TREATMENT PROTOCOLS (Cont.)

Cardiac Arrest (Non-Traumatic)	88
Cold Related Illness	90
Drowning.....	92
Epistaxis	94
Heat Related Illness.....	96
Neonatal Resuscitation	98
Overdose/Poisoning.....	100
Pain Management	102
Respiratory Distress	104
Seizure.....	106
Shock.....	108
Smoke Inhalation	110
Tachycardia/Stable.....	112
Tachycardia/Unstable	114
Ventilation Management	116
OPERATIONS PROTOCOLS	118
Communications	129
Do Not Resuscitate (DNR/POLST).....	121
Documentation	123
Hostile Mass Casualty Incident	124
Inter-Facility Transfer of Patients by Ambulance.....	125
Pediatric Patient Destination	126
Prehospital Death Determination	127
Public Intoxication.....	128
Quality Improvement Review	129
Termination of Resuscitation	130
Transport Destinations.....	131
Trauma Field Triage Criteria.....	132
Waiting Room Criteria.....	135
PROCEDURES PROTOCOLS	136
Cervical Stabilization	137
Continuous Positive Airway Pressure (CPAP).....	138
Electrical Therapy/Defibrillation	139
Electrical Therapy/Synchronized Cardioversion	140
Electrical Therapy/Transcutaneous Pacing	141
Endotracheal Intubation	142
Extraglottic Device	144
First Response Evaluate/Release	145
Hemorrhage Control	146
Medication Administration	147
Needle Cricothyroidotomy.....	149
Needle Thoracentesis.....	150
Tracheostomy Tube Replacement	151

PROCEDURES PROTOCOLS (Cont.)

Traction Splint152
Vagal Maneuvers.....153
Vascular Access154

FORMULARY 155

APPENDICES APP

First Response Low-Risk Alpha Evaluate and Release Form (example) A
Release of Medical AssistanceB
Scope of Practice.....C
Receiving Hospital Directory D
Mass Casualty IncidentE

FOREWORD

EMERGENCY MEDICAL SERVICES PROTOCOL MANUAL

Optimal prehospital care results from a combination of careful patient assessment, essential prehospital emergency medical services, and appropriate medical consultation. The purpose of this manual is to provide guidance for **ALL** prehospital care providers and emergency department physicians within the Clark County EMS System.

The **GOAL** of the manual is to **STANDARDIZE** prehospital patient care in Clark County. It is to be understood that these protocols are guidelines. Nothing contained in these protocols shall be construed to expand the scope of practice of any licensed Attendant beyond that which is identified in the Clark County Emergency Medical Services Regulations and these protocols (Appendix C).

NOTHING contained within these protocols is meant to delay rapid patient transport to a receiving facility. Patient care should be rendered while en-route to a definitive treatment facility.

The General Assessment protocols must be followed in the specific sequence noted. For all other treatment protocols, the algorithm defines the care every patient should receive, usually in the order described.

To maintain the life of a specific patient, it may be necessary, in rare instances, for the physician providing on-line medical consultation, as part of the EMS consultation system, to direct a prehospital provider in rendering care that is not explicitly listed within these protocols, to include administering a patient's own medications which are not part of the approved formulary. To proceed with such an order, both the telemetry physician and the provider must acknowledge and agree that the patient's condition and extraordinary care are not addressed elsewhere within these medical protocols, and that the order is in the best interest of patient care. Additionally, the provider must feel capable, based on the instructions given by the telemetry physician, of correctly performing the directed care. Whenever such care is provided, the telemetry physician and the provider must immediately notify the Office of EMS & Trauma System (OEMSTS) of the extraordinary care situation. In addition, the provider must immediately, upon completion of the call, make available the prehospital care record and documentation specifying the nature of the deviation and the ordering physician's name to the OEMSTS. All such incidents will be entered into the Quality Improvement Review process.

Occasionally a situation may arise in which a physician's order cannot be carried out, e.g., the provider feels the administration of an ordered medication would endanger the patient, a medication is not available, or a physician's order is outside of protocol. If this occurs, the provider must immediately notify the telemetry physician as to the reason the order cannot be carried out, and indicate on the prehospital care record what was ordered, the time, and the reason the order could not be carried out. In addition, the provider must immediately notify the OEMSTS, and upon completion of the call, make available the prehospital care record to the OEMSTS. All such incidents will be entered into the Quality Improvement Review process.

Protocol Key:



Caution / Warning / Alert



Pediatric Treatment Consideration (for patients less than 12 years of age)



Telemetry Contact Required



Specific Protocol



EMT Licensed Attendant and above may perform these steps



AEMT Licensed Attendant and above may perform these steps



Paramedic Licensed Attendant

Definition of a patient:

A patient is any individual that meets at least one of the following criteria:

- 1) A person who has a complaint or mechanism suggestive of potential illness or injury;
- 2) A person who has obvious evidence of illness or injury; or
- 3) A person identified by an informed 2nd or 3rd party caller as requiring evaluation for potential illness or injury.

Pediatric patient considerations:

For patients <18 years old, use the Pediatric Patient Destination protocol.

Pediatric treatment protocols are to be used on children who have not yet experienced puberty. Signs of puberty include chest or underarm hair on males, and any breast development in females.

These protocols have been developed specifically for the Clark County EMS System and represent consensus among all of the Clark County EMS agency medical directors and the District Health Officer. The protocols demonstrate a commitment to a consistent approach to quality patient care.

From time to time, protocols may be added or revised by the District Health Officer upon recommendation by the Medical Advisory Board (MAB). Additional recommendations are welcome and appreciated at any time. They may be submitted to the OEMSTS for consideration and referral to the Medical Advisory Board.

Southern Nevada Health District
Office of Emergency Medical Services & Trauma System
P.O. Box 3902
Las Vegas, Nevada 89127

Physical address:
280 S Decatur Blvd
Las Vegas, NV 89152
Office Hours: Mon-Fri 8:00 am to 4:30 pm

Questions may also be telephoned to EMS staff at (702) 759-1050, or visit our website at <http://www.southernnevadahealthdistrict.org/ems/index.php>.

District Health Officer: Fermin Leguen, MD, MPH

EMS Agency Medical Directors who serve on the Medical Advisory Board:

Dan Rollins, MD, Boulder City Fire Department

Scott Scherr, MD, Clark County Fire Department and Guardian Elite Medical Services

Michael Holtz, MD, Clark County Fire Department

Jessica Leduc, DO, Henderson Fire Department

Nate Jenson, DO, Mesquite Fire & Rescue

Ryan Hodnick, DO, Moapa Valley Fire District and Mt. Charleston Fire Protection District

Kelly Morgan, MD, North Las Vegas Fire Department and Las Vegas Fire and Rescue (Interim)

Mike Barnum, MD, American Medical Response

Jeff Davidson, MD, MedicWest Ambulance

David Obert, DO, Community Ambulance

OEMSTS Staff:

Christian Young, MD, EMS & Trauma System Medical Director [YoungC@snhd.org]

John Hammond, BS, Paramedic, EMS & Trauma System Manager [Hammond@snhd.org]

Laura Palmer, MSML, Paramedic, EMS & Trauma System Supervisor [PalmerL@snhd.org]

Roni Mauro, Paramedic, EMS Field Representative [Mauro@snhd.org]

Scott Wagner, BS, NR Paramedic, EMS Field Representative [Wagner@snhd.org]

Rae Pettie, EMSTS Program/Project Coordinator [Pettie@snhd.org]

Michelle Stanton, Senior Administrative Assistant [Stanton@snhd.org]

Hospitals:

Boulder City Hospital: 901 Adams Blvd, Boulder City, NV 89005 (702) 293-4111

Centennial Hills Hospital: 6900 N Durango Dr, Las Vegas, NV 89149 (702) 629-1210

Desert Springs Hospital Medical Center: 2075 E Flamingo Blvd, Las Vegas, NV 89119 (702) 369-7772

Henderson Hospital: 1050 Galleria Drive, Henderson, NV 89011 (702) 963-7000

Mesa View Regional Hospital: 1299 Bertha Howe Ave, Mesquite, NV 89027 (702) 756-3408

Mike O'Callaghan Federal Medical Center: 4700 N Las Vegas Blvd, Las Vegas, NV 89115 (702) 653-3682

MountainView Hospital: 3100 N Tenaya, Las Vegas, NV 89128 (702) 345-4270

North Vista: 1409 E Lake Mead Blvd North, Las Vegas, NV 89030 (702) 657-5512

Southern Hills Hospital: 9300 W Sunset, Las Vegas, NV 89148 (702) 880-2800

Spring Valley Hospital & Medical Center: 5400 S Rainbow, Las Vegas, NV 89118 (702) 853-3611

St Rose San Martin: 8280 W Warm Springs, Las Vegas, NV 89113 (702) 492-8600

St Rose Siena: 3001 St Rose Pkwy, Henderson, NV 89052 (702) 616-5600

Summerlin Hospital Medical Center: 657 N Town Center Dr, Las Vegas, NV 89144 (702) 233-7000

Sunrise Hospital & Medical Center: 3186 S Maryland Pkwy, Las Vegas, NV 89106 (702) 731-8000

University Medical Center: 1800 W Charleston Las Vegas, NV 89102 (702) 383-2211

Valley Hospital Medical Center: 620 Shadow Ln, Las Vegas, NV 89106 (702) 388-4000


TERMS AND CONVENTIONS

AED	means Automated External Defibrillator
AMPLE	means Allergies; Medications; Prior history; Last meal eaten; Events leading up to injury/illness
AMS	means Altered Mental Status
ASA	means Acetylsalicylic Acid
BG	means Blood Glucose
BP	means Blood Pressure
BVM	means Bag-Valve-Mask
CCC	means Continuous Cardiac Compressions
CHF	means Congestive Heart Failure
COPD	means Chronic Obstructive Pulmonary Disease
CP	means Chest Pain
CPR	means Cardiopulmonary Resuscitation
CVA	means Cardiovascular Accident
DCAP-BTLS	means Deformities; Contusions; Abrasions; Punctures/Penetrations; Burns; Tenderness; Lacerations; Swelling
DKA	means Diabetic Ketoacidosis
ECG	means Electrocardiogram
ETA	means Estimated Time of Arrival
ETT	means Endotracheal Tube
GCS	means Glasgow Coma Scale
GU	means Genitourinary
HEENT	means Head, Ears, Eyes, Nose, Throat
HPI	means History of Present Illness
HR	means Heart Rate
ICP	means Intracranial Pressure
IM	means Intramuscular
IN	means Intranasal
IO	means Intraosseous
IV	means Intravenous
IVP	means Intravenous Push
IVPB	means Intravenous Piggyback
JVD	means Jugular Venous Distention

MAD	means Mucosal Atomizer Device
MI	means Myocardial Infarction
MOI	means Mechanism of Injury
NRB	means Non-rebreather
NS	means Normal Saline
NV	means Nausea/Vomiting
OEMSTS	means Office of Emergency Medical Services & Trauma System
OPQRST	means Onset; Provokes; Quality; Radiates; Severity; Time (used in evaluating localized pain)
PCI	means Percutaneous Coronary Intervention
PCR	means Patient Care Record/Report
PO	means By Mouth
PRN	means As Needed
q	means Every
ROSC	means Return of Spontaneous Circulation
RR	means Respiratory Rate
RUQ	means Right Upper Quadrant
SAMPLE	means Symptoms; Allergies; Medications; Prior history; Last meal eaten; Events leading up to injury/illness
SL	means Sublingual
SOB	means Shortness of Breath
S/P	means Status/Post
SQ	means Subcutaneous
S/S	means Signs/Symptoms
SVT	means Supraventricular Tachycardia
TCAs	means Tricyclic Antidepressants
TFTC	means Trauma Field Triage Criteria
TIA	means Transient Ischemic Attack
TKO/KVO	means To Keep Open/Keep Vein Open
VF	means Ventricular Fibrillation
VT	means Ventricular Tachycardia
VS	means Vital Signs
WPW	means Wolff-Parkinson-White Syndrome

ADULT TREATMENT PROTOCOLS

General Adult Assessment



Pearls

- For all scenes where patient needs exceed available EMS resources, initial assessment and treatment shall be in accordance with an approved triage methodology.
- Correct life-threatening problems as identified.
- If the ability to adequately ventilate a patient cannot be established, the patient must be transported to the nearest emergency department.
- Never withhold oxygen from a patient in respiratory distress.
- Contact with online medical control should be established by radio. Telephone contact may only be used if the call is routed via a recorded phone patch through FAO at 702-382-9007.

Disposition

- Patients sustaining traumatic injuries shall be transported in accordance with the Trauma Field Triage Criteria Protocol.
- Patients sustaining burn injuries shall be transported in accordance with the Burns Protocol.
- Pediatric patients (<18 y/o for transport purposes only) shall be transported in accordance with the Pediatric Destination Protocol.
- Patients with evidence of a stroke shall be transported in accordance with the Stroke (CVA) Protocol.
- Sexual assault victims <13 y/o shall be transported to Sunrise Hospital.
- Sexual assault victims 13 y/o up to 18 y/o shall be transported to Sunrise Hospital or UMC.
- Sexual assault victims 18 y/o and older shall be transported to UMC.
- For sexual assault victims outside a 50-mile radius from the above facilities, the patient shall be transported to the nearest appropriate facility.
- Stable patients shall be transported to the hospital of their choice, if the patient has no preference the patient should be transported to the closest facility.
- For patients outside a 50 mile radius from protocol designated transport destinations, the licensee providing emergency medical care shall transport the patient to the nearest appropriate facility.

Waiting Room Criteria


Upon arrival in the emergency department, if transfer of care has not occurred in accordance with NRS 450B.790, any patient, excluding patients placed on a legal psychiatric hold, meeting *ALL* the following criteria may be placed in the hospital waiting room or other appropriate location:

1. Normal vital signs
 - A. Heart rate 60 - 100
 - B. Respiratory rate 10 - 20
 - C. Systolic BP 100 - 180
 - D. Diastolic BP 60 - 110
 - E. Room air pulse oximetry >94%
 - F. Alert and oriented x 4
2. Did not receive any parenteral medications during EMS transport except a single dose of analgesia and/or an antiemetic.
3. In the judgment of the Paramedic, does not require continuous cardiac monitoring. Note: Any ECG monitoring initiated by a transferring facility may not be discontinued by EMS personnel.
4. Can maintain a sitting position without adverse impact on their medical condition.
5. Is left with a verbal report to hospital personnel.

Internal Disaster

- If a hospital declares an internal disaster, that facility is to be bypassed for all patients except patients in cardiac arrest or in whom the ability to adequately ventilate has not been established.
- Operational exceptions may be initiated in regard to transport to hospitals on internal disaster.

General Adult Trauma Assessment



History

- Time and mechanism of injury
- Damage to structure or vehicle
- Location in structure or vehicle
- Others injured or dead
- Speed and details of MVC
- Restraints/protective equipment
- Past medical history
- Medications

Signs and Symptoms

- Pain, Swelling
- Deformity, lesions, bleeding
- AMS or unconscious
- Hypotension or shock
- Arrest


Differential (life threatening)

- Tension pneumothorax
- Flail chest
- Pericardial tamponade
- Open chest wound
- Hemothorax
- Intra-abdominal bleeding
- Pelvis/femur fracture
- Spine fracture/cord injury
- Head injury
- Extremity fracture
- HEENT (airway obstruction)
- Hypothermia

Pearls

- Recommended exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Back, Neuro.
- Transport destination is based on the Trauma Field Triage Criteria Protocol.
- Transport should not be delayed for procedures; ideally procedures should be performed enroute when possible.
- BVM is an acceptable method of ventilating and managing an airway if pulse oximetry can be maintained $\geq 90\%$.
- Geriatric patients should be evaluated with a high index of suspicion; occult injuries may be present and geriatric patients can decompensate quickly.

Abdominal / Flank Pain, Nausea & Vomiting



History

- Age
- Medical/surgical history
- Onset
- Quality
- Severity
- Fever
- Menstrual history

Signs and Symptoms

- Pain location
- Tenderness
- Nausea
- Vomiting
- Diarrhea
- Dysuria
- Constipation
- Vaginal bleeding/discharge
- Pregnancy


Differential

- Liver (Hepatitis)
- Gastritis
- Gallbladder
- MI
- Pancreatitis
- Kidney stone
- Abdominal aneurysm
- Appendicitis
- Bladder/prostate disorder
- Pelvic (PID, ectopic pregnancy, ovarian cyst)
- Spleen enlargement
- Bowel obstruction
- Gastroenteritis
- Ovarian and testicular torsion

Pearls

- Recommended Exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Back, Extremities, Neuro.
- Neuro disorders or signs of hypoperfusion/shock in the presence of abdominal pain may indicate an aneurysm.
- Document mental status and vital signs prior to administration of antiemetics & pain management.
- Repeat vital signs after each fluid bolus
- In patients ≥ 35 years old consider cardiac origin. Perform a 12-Lead ECG.
- Consider retroperitoneal palpation for kidney pain.
- Abdominal pain in women of childbearing age should be considered pregnancy until proven otherwise.

Allergic Reaction



History

- Onset and location
- Insect sting or bite
- Food allergy/exposure
- Medication allergy/exposure
- New clothing, soap, detergent
- Past history of reactions
- Past medical history
- Medication history

Signs and Symptoms

- Itching or hives
- Coughing/wheezing or respiratory distress
- Throat or chest constriction
- Difficulty swallowing
- Hypotension/shock
- Edema
- Nausea/vomiting

Differential

- Urticarial (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration/airway obstruction
- Asthma/COPD
- CHF

Pearls

- Recommended Exam: Mental Status, Skin, Heart, Lung.
- Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.
- Epinephrine is a first-line drug that should be administered in acute anaphylaxis (moderate / severe symptoms). IM Epinephrine (1:1,000) should be administered in priority before or during attempts at IV or IO access.
- Contact Medical Control for refractory anaphylaxis.
- Consider ETCO₂ monitoring.
- Hypovolemia or distributive shock should be addressed with a fluid bolus prior to the administration of push-dose pressors.
- While there are no absolute contraindications to epinephrine, it should be used with caution in elderly patients, patients with known cardiovascular disease, or significant tachycardia or hypertension, and should be administered only when the patient's signs and symptoms are severe.

Severity

- **Mild** reactions involve skin rashes, itchy sensation, or hives with no respiratory involvement.
- **Moderate** reactions involve skin disorders and may include some respiratory involvement like wheezing, yet the patient still maintains good tidal volume air exchange.
- **Severe** reactions involve skin disorders, respiratory difficulty, and may include hypotension.


Special Considerations

- Always perform ECG monitoring when administering Epinephrine.
- Consider Dopamine for hypotension refractory to administration of Epinephrine.
- Provide oxygen and airway support as needed.

QI Metrics:

- Epinephrine given appropriately.
- Airway assessment documented.

Altered Mental Status / Syncope



History

- Known diabetic, Medic Alert tag
- Drugs or drug paraphernalia
- Report of drug use or toxic ingestion
- Past medical history
- Medications
- History of trauma
- Change in condition
- Changes in feeding or sleep habits

Signs and Symptoms

- Decreased mental status or lethargy
- Changes in baseline mental status
- Bizarre behavior
- Hypoglycemia
- Hyperglycemia
- Irritability


Differential

- Head trauma
- CNS (stroke, tumor, seizure, infection)
- Cardiac (MI, CHF)
- Hypothermia
- Infection
- Thyroid
- Shock (septic, metabolic, traumatic)
- Diabetes
- Toxicological or ingestion
- Acidosis/Alkalosis
- Environmental exposure
- Hypoxia
- Electrolyte abnormality
- Psychiatric disorder

Pearls

- Recommended Exam: Mental Status, HEENT, Skin, Heart, Lung, Abdomen, Back Extremities, Neuro.
- Pay careful attention to the head exam for signs of injury.
- Be aware of AMS as presenting sign of an environmental toxin or Haz-Mat exposure, and protect personal safety and that of other responders.
- Do not let alcohol confuse the clinical picture; alcohol is not commonly a cause of total unresponsiveness to pain.
- If narcotic overdose or hypoglycemia is suspected, administer Narcan 0.4-2mg or Glucose prior to advanced airway procedures.

Behavioral Emergency



History

- Situational crisis
- Psychiatric illness/medications
- Injury to self or threats to others
- Medic Alert tag
- Substance abuse/overdose
- Diabetes

Signs and Symptoms

- Anxiety, agitation, confusion
- Affect change, hallucinations
- Delusional thoughts, bizarre behavior
- Combative, violent
- Expression of suicidal/homicidal thoughts

Differential

- AMS differential
- Alcohol intoxication
- Toxin/substance abuse
- Medication effect or overdose
- Withdrawal syndromes
- Depression
- Bipolar
- Schizophrenia
- Anxiety disorder

Pearls

- Law enforcement assistance should be requested on all calls involving potentially violent patients.
- Under no circumstances are patients to be transported restrained in the prone position.
- Recommended Exam: Mental Status, Skin, Heart, Lung, Neuro.
- Consider all possible medical/trauma causes for behavior.
- Do not irritate the patient with a prolonged exam.
- EMS providers are mandatory reporters in regard to suspected abuse of any vulnerable person.
- Consider ETCO₂ monitoring.

Excited Delirium Syndrome

- Medical emergency-combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent behavior, insensitivity to pain, hyperthermia, and increased strength.
- Potentially life threatening, and associated with the use of physical control measures including restraints, TASER, or similar device.
- Most common in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulants.


Dystonic Reaction

- Condition causing involuntary muscle movements or spasms typically of the face, neck and upper extremities.
- Typically an adverse reaction to drugs such as Haloperidol (may occur with administration).
- When recognized, administer Diphenhydramine 50 mg IM/IV/IO.

S.A.F.E.R.

- Stabilize the situation by containing and lowering the stimuli.
- Assess and acknowledge the crisis.
- Facilitate the identification and activation of resources (chaplain, family, friends or police).
- Encourage patient to use resources and take actions in his/her best interest.
- Recovery or referral – leave patient in care of responsible person or professional, or transport to appropriate facility.

Bradycardia



History

- Past medical history
- Medications
- Pacemaker

Signs and Symptoms

- HR <60/min with hypotension, acute AMS, chest pain, acute CHF, seizures, syncope, or shock secondary to bradycardia
- Respiratory distress

Differential

- Acute myocardial infarction
- Hypoxia
- Pacemaker failure
- Hypothermia
- Sinus bradycardia
- Athletic
- Head injury (elevated ICP) or stroke
- Spinal cord lesion
- AV block
- Overdose


Pearls

- Recommended Exam: Mental Status, HEENT, Heart, Lung, Neuro.
- Bradycardia causing symptoms is typically <50/minute. Rhythm should be interpreted in the context of symptoms and pharmacological treatment given only when symptomatic, otherwise monitor and reassess.
- Identifying signs and symptoms of poor perfusion caused by bradycardia are paramount.
- Do not delay pacing while waiting for IV access.
- Hypoxemia is a common cause of bradycardia; be sure to oxygenate the patient and provide ventilatory support as needed.

QI Metrics

- High degree blocks correctly identified.
- Pacer pads on patient if Atropine given.
- Patient paced if appropriate.

Burns



History

- Type of exposure (heat, gas, chemical)
- Inhalational injury
- Time of injury
- Past medical history & medications
- Other trauma
- Loss of consciousness
- Tetanus/immunization status

Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/distress
- Wheezing
- Singed facial or nasal hair
- Hoarseness or voice changes

Differential


- Superficial (1st degree) – red and painful
- Partial Thickness (2nd degree) – blistering
- Full Thickness (3rd degree) – painless/charred or leathery skin
- Thermal
- Chemical
- Electrical
- Radiation
- Lightning

Pearls

- Burn patients are trauma patients; evaluate for multisystem trauma.
- Assure whatever has caused the burn, is no longer contacting the injury. (Stop the burning process!)
- Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, Neuro.
- Consider early intubation with patients experiencing significant inhalation injuries.
- Potential CO exposure should be treated with 100% oxygen. (For patients in which the primary event is CO inhalation, transport to a hospital equipped with a hyperbaric chamber is indicated [when reasonably accessible].)
- Circumferential burns to extremities are dangerous due to potential vascular compromise secondary to soft tissue swelling. Elevate extremity.
- Burn patients are prone to hypothermia - never apply ice or cool burns; must maintain normal body temperature.
- Consider ETCO₂ monitoring.

Early Intubation Indications

- Signs of Airway Obstruction
- Hoarseness, Stridor, Dysphagia
- Extensive Deep Facial Burns
- Signs of Respiratory Compromise
 - Accessory Muscle Use
 - Inability to Clear Secretions
 - Poor Oxygenation
- Burns in Mouth
- Total BSA ≥ 40%
- Altered Mentation
- Significant Risk of Edema



Patients meeting the following Criteria shall be transported to the closest appropriate Burn Care Center:

1. Second degree burns >10% body surface area (BSA).
2. Any Third degree burns.
3. Burns that involve the face, hands, feet, genitalia, perineum, or major joints.
4. Electrical burns including lightning injury.
5. Chemical burns.
6. Circumferential burns.
7. Inhalation burns.
8. Burn injury with concomitant trauma

Fluid Resuscitation

- Adults 13 years and above 500 ml NS or LR bolus
- Contact Burn Center Medical Direction for additional boluses or drip rates or if it is a prolonged transport.


Pearls (Electrical)

- Do not contact the patient until you are certain the source of the electric shock has been disconnected.
- Attempt to locate contact points, (entry wound where the AC source contacted the patient; an exit at the ground point); both sites will generally be full thickness.
- Cardiac monitor; anticipate ventricular or atrial irregularity to include V-Tach, V-Fib, heart blocks, etc.
- Attempt to identify the nature of the electrical source (AC vs DC), the amount of voltage and the amperage the patient may have been exposed to during the electrical shock.

Pearls (Chemical)

- Certainly 0.9% NaCl Sol'n or Sterile Water is preferred; however if it is not readily available, do not delay; use tap water for flushing the affected area or other immediate water sources. Flush the area as soon as possible with the cleanest, readily available water or saline solution using copious amounts of fluids.

Cardiac Arrest (Non-Traumatic)



History

- Events leading to arrest
- Estimated down time
- Past medical history
- Medications
- Existence of terminal illness

Signs and Symptoms

- Unresponsive
- Apneic
- Pulseless

Differential

- Medical vs. Trauma
- VF vs. Pulseless VT
- Asystole
- PEA
- Primary cardiac event vs. respiratory or drug overdose

Pearls

- For cardiac arrest patients who are pregnant, manual CPR is recommended.
- For cardiac arrest patients who are pregnant, manual displacement of the uterus to the left side is recommended.

Left uterine displacement using 1-handed technique.



Terry L. Vanden Hoek et al. Circulation. 2010;122:S829-S861



Copyright © American Heart Association, Inc. All rights reserved.



Terry L. Vanden Hoek et al. Circulation. 2010;122:S829-S861




Copyright © American Heart Association, Inc. All rights reserved.

- Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.
- Consider early IO placement if IV is difficult.
- DO NOT HYPERVENTILATE.
- Reassess and document ETT placement using auscultation and ETCO₂ capnography.
- Switch compressors every two minutes.
- Try to maintain patient modesty.
- Mechanical chest compression devices should be used if available in order to provide for consistent uninterrupted chest compressions and crew safety. As noted above, mechanical chest compression devices are not recommended for the pregnant patient.

H's & T's (reversible causes)

- Hypovolemia – Volume infusion
- Hypoxia – Oxygenation & ventilation, CPR
- Hydrogen ion (acidosis) – Ventilation, CPR
- Hypokalemia
- Hyperkalemia -Calcium chloride, sodium bicarbonate, albuterol
- Hypothermia - Warming
- Tension pneumothorax – Needle decompression
- Tamponade, cardiac – Volume infusion
- Toxins – Agent specific antidote
- Thrombosis, pulmonary – Volume infusion
- Thrombosis, coronary – Emergent PCI

Chest Pain (Non Traumatic) and Suspected Acute Coronary Syndrome



History

- Age
- Medications: Viagra, Levitra, Cialis
- Past medical history of MI, angina, diabetes
- Allergies
- Recent physical exertion
- Palliation, provocation
- Quality
- Region, radiation, referred
- Severity (1-10)
- Time of onset, duration, repetition

Signs and Symptoms

- CP, pressure, ache, vise-like pain, tight
- Location, substernal, epigastric, arm, jaw, neck, shoulder
- Radiation of pain
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness
- Time of onset

Differential

- Trauma versus medical
- Anginal versus MI
- Pericarditis
- Pulmonary embolism
- Asthma, COPD
- Pneumothorax
- Aortic dissection or aneurysm
- GE reflux or hiatal hernia
- Esophageal spasm
- Chest injury or pain
- Pleural pain
- Drug overdose (cocaine, methamphetamine)


Pearls

- Recommended exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Diabetics, geriatrics, and female patients often have atypical pain. Have a high index of suspicion.
- Perform a 12-Lead ECG on all patients 35 years old or older experiencing vague jaw/ chest/ abdominal discomfort.
- Perform a 12-Lead ECG within 5 minutes of patient contact.
- The administration of nitroglycerin is contraindicated for any patient who has used erectile dysfunction medications within the last 48 hours.
- Nitroglycerin is contraindicated in any patient with hypotension, bradycardia, or tachycardia in the absence of heart failure and evidence of a right ventricular infarction.

QI Metrics

- 12-Lead ECG within 5 minutes of patient contact.
- Pain reassessed after every intervention.
- Pain control documented.

Childbirth / Labor



History

- Due date
- Time contractions started/ duration/frequency
- Rupture of membranes (meconium)
- Time and amount of any vaginal bleeding
- Sensation of fetal movement
- Pre-natal care
- Past medical and delivery history
- Medications
- Gravida/Para status
- High risk pregnancy

Signs and Symptoms

- Spasmodic pain
- Vaginal discharge or bleeding
- Crowning or urge to push
- Meconium

Differential


- Abnormal presentation (breech, limb)
- Prolapsed cord
- Placenta previa
- Abruptio placenta

Pearls

- Recommended exam (of mother): Mental Status, Heart, Lungs, Abdomen, Neuro.
- Document all times (delivery, contraction duration and frequency).
- Some bleeding is normal; copious amounts of blood or free bleeding is abnormal.
- Record APGAR at one and five minutes after birth.
- APGAR of 7-10 is normal, while 4-7 requires resuscitative measures.

APGAR	Score=0	Score=1	Score=2
• <u>Activity/Muscle Tone</u>	Absent	Arms/legs flexed	Active movement
• <u>Pulse</u>	Absent	Below 100	Above 100
• <u>Grimace/Reflex Irritability</u>	No response	Grimace	Sneeze, cough, pulls away
• <u>Appearance/Skin Color</u>	Blue-Grey, pale all over	Normal, except extremities	Normal over entire body
• <u>Respiration</u>	Absent	Slow, irregular	Good, crying

Cold-Related Illness



History

- Age, very young and old
- Exposure to decreased temperatures, but may occur in normal temperatures
- Past medical history/medications
- Drug or alcohol use
- Infections/sepsis
- Time of exposure/wetness/wind chill

Signs and Symptoms

- AMS/coma
- Cold, clammy
- Shivering
- Extremity pain
- Bradycardia
- Hypotension or shock

Differential

- Sepsis
- Environmental exposure
- Hypoglycemia
- Stroke
- Head injury
- Spinal cord injury

Pearls

- Recommended exam: Mental Status, Heart, Lung, Abdomen, Extremities, Neuro.
- Extremes of age are more prone to cold emergencies.
- Obtain and document patient temperature.
- If temperature is unknown, treat the patient based on suspected temperature.
- Active warming includes hot packs that can be used on the armpit and groin; care should be taken not to place the packs directly on the skin.
- Warm saline or lactated ringers IV may be used.
- Recognize the cardiac arrest resuscitation guidelines for the hypothermic patient.

Hypothermia Categories

- Mild 90° - 95° F (33° - 35° C)
- Moderate 82° - 90° F (28° - 32° C)
- Severe <82 degrees F (<28° C)


Hypothermia Mechanisms

- Radiation
- Convection
- Conduction
- Evaporation

Active Heating Measures

- Hot packs to the armpits and groin (do not place directly onto skin)

Drowning



CAVEATS:

1. Adequate ventilation is KEY!!!
2. For patients breathing on their own, start Oxygen 15 L NRB; for patients not adequately breathing → BVM
3. Do not suction foam in airway, just bag through it initially.
4. For drowning victims in cardiac arrest, emphasis should be on good oxygenation/ventilation → use traditional 30:2 CPR (no continuous compressions).

History

- Submersion in fluid, regardless of depth
- Possible history of trauma (dive)
- Duration of immersion
- Temperature of water or possibility of hypothermia
- Degree of water contamination

Signs and Symptoms

- Unresponsive
- Mental status changes
- Decreased or absent vital signs
- Vomiting
- Coughing, wheezing, rales, stridor, rhonchi
- Apnea
- Frothy/foamy sputum

Differential

- Trauma
- Pre-existing medical condition
- Barotrauma
- Decompression illness
- Post-immersion syndrome


Pearls

- Recommended Exam: Trauma Survey, Head, Neck, Chest, Abdomen, Back, Extremities, Skin, Neuro.
- Ensure scene safety.
- Hypothermia is often associated with submersion incidents.
- All patients should be transported for evaluation because of potential for worsening over the next several hours.

QI Metrics

- Submit the SNHD Submersion Incident Report Form.

Epistaxis



History

- Age
- Past Medical History
- Medications (HTN, Anticoagulants, aspirin, NSAIDS)
- Previous episodes of epistaxis
- Trauma
- Duration of bleeding
- Quantity of bleeding

Signs and Symptoms

- Bleeding from nasal passages
- Pain
- Nausea
- Vomiting


Differential

- Trauma
- Infection (viral URI or Sinusitis)
- Allergic rhinitis
- Lesions (polyps, ulcers)
- Hypertension

Pearls

- Recommended exam: Mental Status, HEENT, Lungs, Neuro
- It is very difficult to quantify the amount of blood loss with epistaxis
- Bleeding may be also occurring posteriorly. Evaluate for posterior blood loss by examining the posterior pharynx.
- Anticoagulants include warfarin (Coumadin), heparin, enoxaparin (Lovenox), dabigatran (Pradaxa), rivaroxaban (Xarelto), and many other over the counter headache relief powders.
- Anti-platelet agents like aspirin, clopidogrel (Plavix), aspirin/dipyridamole (Aggrenox), and ticlopidine (Ticlid) can contribute to bleeding.

Heat-Related Illness



History

- Age, very old and young
- Exposure to increased temperatures and/or humidity
- Past medical history/medications
- Time and duration of exposure
- Poor PO intake, extreme exertion
- Fatigue and/or muscle cramping

Signs and Symptoms

- AMS/coma
- Hot, dry, or sweaty skin
- Hypotension or shock
- Seizures
- Nausea

Differential

- Fever
- Dehydration
- Medications
- Hyperthyroidism
- DTs
- Heat cramps, heat exhaustion, heat stroke
- CNS lesions or tumors

Pearls

- Recommended exam: Mental Status, Skin, Heart, Lung, Abdomen, Extremities, Neuro.
- Extremes of age are more prone to heat emergencies.
- Cocaine, amphetamines, and salicylates may elevate body temperatures.
- Sweating generally disappears as body temperatures rise over 104° F (40° C).
- Intense shivering may occur as patient is cooled.
- Active cooling includes application of cold packs or ice (not directly on skin), fanning either by air conditioning or fanning.
- Cold Saline is not to be administered for the treatment of hyperthermia unless directed by telemetry physician.

Heat Cramps

- Consist of benign muscle cramping caused by dehydration and is not associated with an elevated temperature.

Heat Exhaustion

- Consists of dehydration, salt depletion, dizziness, fever, AMS, headache, cramping, N/V. Vital signs usually consist of tachycardia, hypotension and elevated temperature.


Heat Stroke

- Consists of dehydration, tachycardia, hypotension, temperature >104° F (40° C), and AMS.

Active Cooling Measures

- Cold packs
- Ice (do not place directly onto patient's skin)
- Fanning
- Air Conditioning

Hyperkalemia (Suspected)



History

- History of renal failure
- History of dialysis
- Trauma, crush injury

Signs and Symptoms

- Cardiac conduction disturbances
- Irritability
- Abdominal distension
- Nausea
- Diarrhea
- Oliguria
- Weakness


Differential

- Cardiac disease
- Renal failure
- Dialysis
- Trauma

Pearls

- **Patients must have suspected hyperkalemia *OR* electrocardiographic findings consistent with hyperkalemia (bradycardia with widening QRS complexes) BEFORE initiating treatment.**
- **Calcium Chloride is contraindicated in patients taking digitalis products.**
- **Hyperkalemia is defined as a potassium level higher than 5.5 mmol/L.**
- **Potassium of 5.5 - 6.5 mmol/L - Tall tented T waves.**
- **Potassium of 6.5 - 7.5 mmol/L - Loss of P waves.**
- **Potassium of 7.5 - 8.5 mmol/L - Widening QRS.**
- **Potassium of >8.5 mmol/L - QRS continues to widen, approaching sine wave.**

Obstetrical Emergency



History

- Medical history
- Hypertension medication
- Prenatal care
- Prior pregnancies/births
- Previous pregnancy complications

Signs and Symptoms

- Vaginal bleeding
- Abdominal pain
- Seizures
- Hypertension
- Severe headache
- Visual changes
- Edema of the hands or face


Differential

- Pre-eclampsia/eclampsia
- Placenta previa
- Placenta abruptio
- Spontaneous abortion

Pearls

- Recommended exam: Mental Status, Heart, Lung, Abdomen, Neuro.
- Severe headache, vision changes or RUQ pain may indicate pre-eclampsia.
- In the setting of pregnancy hypertension is defined as >140 systolic or >90 diastolic or a relative increase of 30 systolic and 20 diastolic from the patient's normal pre-pregnancy BP.
- Maintain left lateral position.
- Ask patient to quantify bleeding - number of pads used per hour.
- Any pregnant patient involved in a MVC should be seen by a physician for evaluation.
- Postpartum Eclampsia/Pre-Eclampsia commonly presents up to 48 hours after childbirth. If symptomatic, treat as Eclampsia/Pre-Eclampsia.
- May present up to 6 weeks after childbirth, Assess for history or Pre-Eclampsia/Eclampsia during pregnancy or delivery.

Overdose/Poisoning



History

- Ingestion or suspected ingestion of a potentially toxic agent
- Substance ingested, route, quantity
- Time of ingestion
- Reason (suicidal, accidental, criminal)
- Available medications in home
- Past medical history, medications

Signs and Symptoms

- Mental status changes
- Hypotension/hypertension
- Decreased respiratory rate
- Tachycardia, dysrhythmias
- Seizures
- SLUDGE
- Malaise, weakness
- GI symptoms
- Dizziness
- Syncope
- Chest pain

Differential

- TCA overdose
- Acetaminophen OD
- Aspirin
- Depressants
- Stimulants
- Anticholinergic
- Cardiac medications
- Solvents, alcohols, cleaning agents, insecticides


Pearls

- Recommended exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Neuro.
- Calcium Chloride is contraindicated in patients taking digitalis products.
- Overdose or toxin patients with significant ingestion/exposure should be closely monitored and aggressively treated. Do not hesitate to contact medical control if needed.
- In the case of cyanide poisoning, altered mental status may be profound. Profound altered mental status can be defined as a deficit that includes disorientation, bewilderment and difficulty following commands.
- If patient is suspected to have narcotic overdose/hypoglycemia, administer Narcan/ Glucose prior to extraglottic device/intubation.
- Poison Control: 1-800-222-1222

Agents

- Acetaminophen: Initially normal or N/V. Tachypnea and AMS may occur later. Renal dysfunction, liver failure and/or cerebral edema may manifest.
- Depressants: Decreased HR, BP, temp and RR.
- Anticholinergic: Increased HR, increased temperature, dilated pupils and AMS changes.
- Insecticides: May include S/S of organophosphate poisoning.
- Solvents: N/V, cough, AMS.
- Stimulants: Increased HR, BP, temperature, dilated pupils, seizures, and possible violence.
- TCA: Decreased mental status, dysrhythmias, seizures, hypotension, coma, death.

Pain Management



History

- Age
- Location, duration
- Severity (1-10)
- Past medical history
- Pregnancy status
- Drug allergies and medications

Signs and Symptoms

- Severity (pain scale)
- Quality
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

Differential

- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural, respiratory
- Neurogenic
- Renal (colic)


Pearls

- Recommended exam: Respiratory Status, Mental Status, Area of pain, Neuro.
- Pain severity (1-10) is a vital sign to be recorded before and after medication administration and patient hand off.
- Monitor BP and respirations closely as sedative and pain control agents may cause hypotension and/or respiratory depression.
- Consider patient's age, weight, clinical condition, use of drugs/alcohol, exposure to opiates when determining initial opiate dosing. Weight based dosing may provide a standard means of dosing calculation, but it does not predict response.
- Exercise care when administering opiates and benzodiazepines; this combination results in deeper anesthesia with significant risk of respiratory compromise.
- Burn patients may require more aggressive dosing.
- Administration of Droperidol can result in hypotension, QT prolongation and Torsades de Pointes.

QI Metrics

- Vital signs with O₂ sats recorded.
- Pain scale documented before and after intervention.
- Vital signs repeated after intervention.
- If considering repeat administration of pain medications, nasal cannula capnography must be utilized.

Pulmonary Edema/CHF



History

- Congestive heart failure
- Past medical history
- Medications
- Cardiac history

Signs and Symptoms

- Respiratory distress, bilateral rales
- Apprehension, orthopnea
- JVD
- Pink, frothy sputum
- Peripheral edema
- Diaphoresis
- Hypotension, shock
- Chest pain

Differential

- MI
- Congestive heart failure
- Asthma
- Anaphylaxis
- Aspiration
- COPD
- Pleural effusion
- Pneumonia
- Pericardial tamponade
- Toxic exposure


Pearls

- The administration of nitroglycerin is contraindicated for any patient who has used erectile dysfunction medications within the last 48 hours.
- Carefully monitor the patient as you administer interventions.
- Consider MI.
- Allow patient to maintain position of comfort.
- Consider dose related effects of Dopamine: 2-10 mcg/kg/min increases myocardial contractility and HR, improves BP via vasoconstriction; 10-20 mcg/kg/min causes vasoconstriction of renal, mesenteric, and peripheral blood vessels that can result in poor perfusion and renal failure.

QI Metrics

- Blood pressure reassessed after each nitroglycerin dose.
- ETCO₂ monitored.

Respiratory Distress



History

- Asthma, COPD, CHF, chronic bronchitis, emphysema
- Home treatment (oxygen, nebulizers)
- Medication
- Toxic exposure

Signs and Symptoms

- Shortness of breath
- Pursed lip breathing
- Decreased ability to speak
- Increased respiratory rate and effort
- Wheezing, rhonchi
- Use of accessory muscles
- Fever, cough
- Tachycardia


Differential

- Asthma
- Anaphylaxis
- Aspiration
- COPD
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pneumothorax
- Cardiac (MI or CHF)
- Pericardial tamponade
- Hyperventilation
- Inhaled toxin

Pearls

- Recommended exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro.
- Pulse oximetry and end tidal continuous waveform capnography must be monitored.
- Consider MI.
- Allow the patient to assume a position of comfort.

Seizure



History

- Reported or witnessed seizure activity
- Previous seizure history
- Seizure medications
- History of trauma
- History of diabetes
- History of pregnancy
- Time of seizure onset
- Number of seizures
- Alcohol use, abuse, or abrupt cessation
- Fever

Signs and Symptoms

- Decreased mental status
- Sleepiness
- Incontinence
- Observed seizure activity
- Evidence of trauma
- Unconsciousness

Differential

- CNS trauma
- Tumor
- Metabolic, hepatic or renal failure
- Hypoxia
- Electrolyte abnormality (Na, Ca, Mg)
- Drugs, medication non-compliance
- Infection, fever
- Alcohol withdrawal
- Eclampsia
- Stroke
- Hyperthermia
- Hypothermia

Pearls

- Recommended exam: Mental Status, HEENT, Heart, Lungs, Extremities, Neuro.
- Benzodiazepines are effective in terminating seizures; do not delay IM/IN administration while initiating an IV.
- Status epilepticus is defined as two or more seizures successively without an intervening lucid period, or a seizure lasting over five minutes.
- Grand mal seizures (generalized) are associated with loss of consciousness, incontinence and oral trauma.
- Focal seizures affect only part of the body and are not usually associated with a loss of consciousness.
- Be prepared to address airway issues and support ventilations as needed.
- Consider ETCO₂ monitoring.

Sepsis (Suspected)

 **General Adult Assessment**

- E** Oxygen Keep SpO2 >94%
- A** Vascular Access
- P** Cardiac Monitoring/Capnography

Does the patient have a known or suspected source of infection AND TWO of the following criteria?

- SBP <90 mm Hg
- HR >90/min
- Respiratory rate >20
- Altered mental status
- Temperature > or = 100.4 F or < or = 96.8 F
- Persistent EtCo2 <25 on waveform capnography

YES

NO

Facility Code Sepsis notification telemetry immediately upon recognition of patient meeting Sepsis criteria

Alternate appropriate treatment protocols as indicated

A	NS or LR bolus 500 ml IV/IO, reassess criteria and re-examine. May repeat x3 for SBP <90, with no rales on lung exam. Max dose 2 liters.
P	If SBP <90 after 2 liters IVF PUSH DOSE EPINEPHRINE 1:100,000 5.0 mcg-10.0 mcg IV/IO, may repeat q 2-5 min to maintain SBP >90 (0.5 ml-1 ml of a 1:100,000 solution) OR
	PUSH DOSE PHENYLEPHRINE 100 mcg – 200 mcg IV/IO, may repeat q 2-5 min to maintain SBP >90 (1 ml – 2 ml of a 100 mcg/ml solution)
	Consider DOPAMINE 5-20 mcg/kg/min; titrate to SBP >90

Continue General Adult Assessment

History

- Age (Common in elderly and very young)
- Presence and duration of fever
- Previously documented infection or illness (UTI, Pneumonia, meningitis, encephalitis, cellulitis, abscesses, etc)
- Recent surgery or invasive procedure
- Any recent hospitalization
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Bedridden or immobile patients
- Prosthetic or indwelling devices
- Immunization status
- Open wounds, even minor ones

Signs and Symptoms

- Hyper or hypothermia
- Rash and/or excessive bruising
- Chills
- Myalgia
- Markedly decreased urine output
- Altered mentation
- Delayed capillary refill
- Elevated blood glucose (unless diabetic)

Differential

- **Cardiogenic Shock**
- Hypovolemic Shock
- Dehydration
- Hyperthyroidism
- Medication/drug interaction
- Non-septic infection
- Allergic reaction/anaphylaxis
- Toxicological emergency

Pearls

- Early recognition of Sepsis allows for attentive care and early administration of antibiotics.
- Aggressive IV fluid therapy is the most important prehospital treatment for sepsis. Suspected septic patients should receive repeated fluid boluses (to a max total of 2 liters) while being checked frequently for signs of pulmonary edema, especially those patients with known history of CHF or ESRD on dialysis. STOP fluid resuscitation in the setting of pulmonary edema.
- Time IVF bolus was initiated and total amount given is to be recorded and reported to hospital staff at patient hand off.
- Septic patients are especially susceptible to traumatic lung injury and ARDS. If artificial ventilation is necessary, avoid ventilating with excessive tidal volumes. If CPAP is utilized, airway pressure should be limited to 5 cm H₂O.
- Attempt to identify source of infection (skin, respiratory, etc.) and relay previous treatments and related history to the ED physician and nursing staff.
- Elevated serum lactate levels are a useful marker of hypoperfusion in sepsis and often become elevated prior to the onset of hypotension. ETCO₂ levels are inversely proportional to serum lactate levels.
- Disseminated intravascular Coagulation (DIC) is an ominous, late stage manifestation of sepsis characterized by frank, extensive bruising, bleeding from multiple sites, and finally tissue death.
- Conditions such as Crohn's, psoriasis, rheumatoid arthritis and other autoimmune disorders are now being treated with medications that impair the immune system. These patients need to be considered as immunocompromised.
- Hypovolemia or distributive shock should be addressed with a fluid bolus prior to the administration of push-dose pressors.
- While there are no absolute contraindications to epinephrine, it should be used with caution in elderly patients, patients with known cardiovascular disease, or significant tachycardia or hypertension, and should be administered only when the patient's signs and symptoms are severe.

QI Metrics

- Vital signs to include blood pressure, heart rate, respiratory rate, SpO₂, and EtCO₂ documented throughout transport.
- Vital signs before, during, and after IVF administration.
- Documentation of the time IVF was started and total amount given.

Shock




For patients with known adrenal insufficiency, administer patient's own Solu-Cortef (hydrocortisone) as prescribed

General Adult Assessment

- E** Oxygen Keep SpO₂ >94%
- A** Vascular Access
- P** Cardiac monitor/capnography

Alternative appropriate treatment protocols as indicated



General Trauma

A NS or LR bolus 1000 ml IV/IO; may repeat x 1 with no rales on lung exam

Appropriate arrhythmia protocol as indicated

P 12-Lead ECG
Obtain waveform capnography

A NS or LR bolus 500 ml IV/IO; if no rales on lung exam, may repeat x 1

P

PUSH DOSE EPINEPHRINE
1:100,000
5.0mcg – 10 mcg IV/IO, may repeat q 2-5 min to maintain SBP >90 (0.5 ml-1 ml of a 1:100,000 solution)
OR
PUSH DOSE PHENYLEPHRINE
100 mcg – 200 mcg IV/IO q 2-5 min to maintain SBP >90 (1 ml – 2ml of 100 mcg/ml solution)
OR
DOPAMINE
5-20 mcg/kg/min, titrate to SBP>90

P

PUSH DOSE PHENYLEPHRINE
100 mcg – 200 mcg IV/IO q 5 min to maintain SBP>90 (1 ml- 2 ml of a 100mcg/ml solution)
OR
DOPAMINE
5-20 mcg/kg/min; titrate to SBP>90

Obtain waveform capnography

Obtain waveform capnography

Continue General Adult Assessment

History

- Blood loss-vaginal bleeding, ectopic, GI bleeding or AAA
- Fluid loss-vomiting, diarrhea, fever
- Infection
- Cardiac tamponade
- Medications
- Allergic reaction
- Pregnancy
- History of poor oral intake

Signs and Symptoms

- Restlessness, confusion
- Weakness, dizziness
- Weak rapid pulse
- Pale, cool, clammy skin
- Delayed capillary refill
- Hypotension
- Coffee-ground emesis
- Tarry stools

Differential

- Hypovolemic shock
- Cardiogenic shock
- Septic shock
- Neurogenic shock
- Anaphylactic shock
- Ectopic pregnancy
- Dysrhythmias
- Pulmonary embolism
- Tension pneumothorax
- Medication effect or overdose
- Vasovagal
- Physiologic (pregnancy)

Pearls

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Hypotension can be defined as a systolic BP of <90. This is not always reliable and should be interpreted in context and patient's typical BP, if known. Shock may present with a normal BP initially.
- Hypovolemia or distributive shock should be addressed with a fluid bolus prior to the administration of push-dose pressors.
- While there are no absolute contraindications to epinephrine, it should be used with caution in elderly patients, patients with known cardiovascular disease, or significant tachycardia or hypertension, and should be administered only when the patient's signs and symptoms are severe.
- Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the only manifestation.
- Consider all possible causes of shock and treat per appropriate protocol.
- An ETCO₂ measurement of <25 mm/hg is indicative of shock

Hypovolemic shock

- Hemorrhage, trauma, GI bleeding, ruptured aortic aneurysm, or pregnancy related bleeding

Cardiogenic shock

- Heart failure, MI, cardiomyopathy, myocardial contusion, toxins

Distributive shock

- Sepsis (consider telemetry of code sepsis to receiving facility), anaphylaxis, neurogenic, toxins

Obstructive shock

- Pericardial tamponade, pulmonary embolus, tension pneumothorax

For patients with known adrenal insufficiency, administer patient's own Solu-Cortef (hydrocortisone) as prescribed.

Causes of Adrenal Insufficiency:


Addison's Disease

Congenital Adrenal Hyperplasia

Long term administration of steroids

Others

Smoke Inhalation



History

- Exposed to smoke in a structure fire
- Exposed to smoke in a vehicle fire
- Exposed to smoke from other sources, industrial, confined space, wilderness fire, etc.

Signs and Symptoms

- Facial burns
- Singed nasal hairs or facial hair
- Shortness of breath
- Facial edema
- Stridor
- Grunting respirations

Differential

- COPD
- CHF
- Toxic inhalation injury
- Caustic inhalation injury

Pearls

- Protect yourself and your crew.
- Have a high index of suspicion when treating patients at the scene of a fire.
- If the medication is not available on scene do not delay transport waiting for it.
- Carefully monitor respiratory effort and correct life threats immediately.
- Decide early on if you want to intubate as burned airways swell, making intubation difficult.
- Profound altered mental status can be defined as a deficit that includes disorientation, bewilderment and difficulty following commands.

Preparation and Administration of Hydroxocobalamin


Complete Starting Dose: 5 g

1. Reconstitute: Place the vial in an upright position. Add **200 mL** of 0.9% Sodium Chloride Injection* to the vial using the transfer spike. **Fill to the line.**


* 0.9% Sodium Chloride Injection is the recommended diluent (diluent not included in the kit). Lactated Ringer's Solution and 5% Dextrose Injection have also been found to be compatible with Hydroxocobalamin.

2. Mix: The vial should be repeatedly inverted or rocked, **NOT** shaken, for at least **60 seconds** prior to infusion.

3. Infuse Vial: Use vented intravenous tubing, hang and infuse over **15 minutes**.



STEMI (Suspected)



History

- Age
- Medication: Viagra, Levitra, Cialis
- Past Medical History of MI, angina, diabetes
- Allergies
- Recent Physical Exertion
- Palpitation, provocation
- Quality
- Region, radiation, referred
- Severity (1-10)
- Time of onset, duration, repetition

Signs and Symptoms

- CP, pressure, ache, vice-like pain, tight
- Location, substernal, epigastric, arm, jaw, neck, shoulder
- Radiation of pain
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness
- Time of onset

Differential

- Trauma versus medical
- Anginal versus MI
- Pericarditis
- Pulmonary embolism
- Asthma, COPD
- Pneumothorax
- Aortic dissection or aneurysm
- GE reflux or hiatal hernia
- Esophageal spasm
- Chest injury or pain
- Pleural pain
- Drug overdose (cocaine, methamphetamines)


Pearls

- Recommended exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Diabetics, geriatrics, and female patients often have atypical pain. Have a high index of suspicion.
- Perform a 12-Lead ECG on all patients 35 years old and older experiencing vague jaw/ chest/ abdominal discomfort.
- Perform a 12-Lead ECG within 5 minutes of patient contact.
- The administration of nitroglycerin is contraindicated for any patient who has used erectile dysfunction medications within the last 48 hours.

QI Metrics

- 12-Lead ECG within 5 minutes of patient contact.
- Pain reassessed with every intervention.
- Pain control documented.

Stroke (CVA)



History

- Previous CVA, TIAs
- Previous cardiac/vascular surgery
- Associated diseases: diabetes, HTN
- CAD
- Atrial Fibrillation
- Medications
- History of trauma

Signs and Symptoms

- AMS
- Weakness, paralysis
- Blindness or other sensory loss
- Aphasia, dysarthria
- Syncope
- Vertigo, dizziness
- Vomiting
- Headache
- Seizures
- Respiratory pattern change
- Hypertension, hypotension

Differential

- AMS
- TIA
- Seizure
- Hypoglycemia
- Tumor
- Trauma
- Dialysis/ Renal Failure

Pearls

- Recommended exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro.
- Determine time of onset of symptoms or last time patient was seen normal
- Transport to an appropriate Stroke Center or Endovascular Treatment Center

Stroke Centers

- Centennial Hills
- Desert Springs
- Henderson
- MountainView
- Southern Hills
- Spring Valley
- St Rose Siena
- St Rose San Martin
- Summerlin
- Sunrise
- UMC
- Valley

NIR Capable Centers

- Centennial Hills
- Henderson Hospital
- Southern Hills Hospital
- Spring Valley
- St Rose Siena
- Sunrise
- UMC
- Valley

QI Metrics

- Complete the RACE assessment in less than five minutes
- Time of symptom onset documented
- Blood glucose documented
- 12-Lead EKG completed
- Scene time <10 minutes
- Telemetry to receiving facility

Rapid Arterial Occlusion Evaluation (RACE) Scale

An EMS Assessment Tool for Acute Ischemic Stroke

(Sensitivity 85%, Specificity 68%)

Test Item	Score = 0	Score = 1	Score = 2	Patient Score
Facial Palsy	Absent	Mild	Moderate/Severe	
Arm Motor	Normal/Mild	Moderate	Severe	
Leg Motor	Normal/Mild	Moderate	Severe	
Head/Gaze Deviation	Absent	Present	N/A	
Aphasia* (if righthemiparesis)	Performs Both Tasks	Performs 1 Task	Performs Neither Tasks	
Agnosia* (if lefthemiparesis)	Patient Recognizes Arm and Impairment	Unable to Recognize Arm or Impairment	Unable to Recognize BOTH Arm and Impairment	
			TOTAL SCORE = (0-9)	

*Aphasia: Ask the patient to: 1. "Close your Eyes" AND 2. "Make a Fist"

*Agnosia: Ask the patient and evaluate recognition of deficit:


1. While showing paretic arm: "Whose arm is this?"
2. Ask patient: "Can you lift both arms and clap?"

If RACE Score = 5 or greater, patient may have an ischemic stroke with a large vessel occlusion

Reference:

Natalia Pérez de la Ossa, et al. (2014). Design and Validation of a Prehospital Stroke Scale to Predict Large Arterial Occlusion: The Rapid Arterial Occlusion Evaluation Scale. *Stroke*, 45, 87-91. Retrieved from <http://stroke.ahajournals.org/content/45/1/87.full>

Tachycardia/Stable (Normal Mental Status, Palpable Radial Pulse)



History

- Medications (aminophylline, diet pills, thyroid supplements, decongestants, digoxin)
- Diet (caffeine)
- Drugs (cocaine, methamphetamines)
- Past medical history
- Syncope/near syncope
- History of palpitations/racing heart

Signs and Symptoms

- Heart rate >150
- Dizziness, CP, SOB
- Diaphoresis
- CHF


Differential

- Heart disease (WPW, valvular)
- Sick sinus syndrome
- MI
- Electrolyte imbalance
- Exertion, fever, pain, emotional stress
- Hypoxia
- Hypovolemia
- Drug effect, overdose
- Hyperthyroidism

Pearls

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Carefully monitor patients as you treat them; stable tachycardia may convert to unstable rhythms/conditions quickly.
- Sedate patients prior to cardioversion, if time allows.

Tachycardia / Unstable (Mental Status Changes, No Palpable Radial Pulse)



History

- Medications (aminophylline, diet pills, thyroid supplements, decongestants, digoxin)
- Diet (caffeine)
- Drugs (cocaine, methamphetamines)
- Past medical history
- Syncope/near syncope
- History of palpitations/racing heart

Signs and Symptoms

- Cardiac arrest
- Heart rate >150
- Dizziness, CP, SOB
- Diaphoresis
- CHF

Differential

- Heart disease (WPW, valvular)
- Sick sinus syndrome
- MI
- Electrolyte imbalance
- Exertion, fever, pain, emotional stress
- Hypoxia
- Hypovolemia
- Drug effect, overdose
- Hyperthyroidism

Pearls

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- If patient is in arrest, efforts should focus on quality chest compressions and rhythm correction.
- Administer Adenosine at a proximal IV site, rapidly followed by a saline flush.

Ventilation Management

Use supplemental oxygen to maintain an oxygen saturation of >94% or >90% for patients on home oxygen for chronic conditions

E	Basic Airway Maneuvers -Open Airway Chin Lift/Jaw Thrust -NPA or OPA as needed -Suction as needed	Respiratory Distress and/or Tracheostomy Tube Replacement Protocol if needed
	Consider Cervical Stabilization	
	Consider Altered Mental Status/Syncope	

E	Administer oxygen
	BVM as needed

Intervention effective?

A	Extraglottic Airway	Obtain Vascular Access
	Endotracheal Intubation	ECG Monitor
P	For Nasotracheal Intubation Prep the Nostrils With PHENYLEPHRINE 1-2 Sprays Each Or OXYMETAZOLINE 1-2 sprays each nostril and LIDOCAINE 2% Lubricant	
	Consider Induction Administer ETOMIDATE 0.3 mg/kg IV/IO, max dose 30 mg OR; KETAMINE 2 mg/kg IV/IO Or 4 mg/kg IM	
	Post-Intubation Sedation Administer MIDAZOLAM 0.1 mg/kg IV/IN/IM/IO; max dose 5mg May Repeat X 1 after 5 min at 0.05 mg/kg IV/IN/IM/IO, max dose 2.5 mg, further doses with Physician Order OR; DIAZEPAM 5 mg IV/IO; May Repeat Dose after 5 Minutes With Physician Order	
	Post-Sedation Analgesia Administer FENTANYL 1 mcg/kg; max dose 100 mcg.	

Extraglottic/ETT placement successful?

Able to ventilate without extraglottic/ETT in place?

Cricothyroidotomy

Continue Care and Transport as Appropriate

Always weigh the risks and benefits of endotracheal intubation in the field against transport. All prehospital endotracheal intubations are considered high risk. If ventilation/oxygenation is adequate, transport may be the best option. The most important airway device and the most difficult to use correctly and effectively is the Bag Valve Mask (not the laryngoscope). Few prehospital airway emergencies cannot be temporized or managed with proper BVM techniques.

DIFFICULT AIRWAY ASSESSMENT:

Difficult BVM Ventilation-MOANS: Difficult **M**ask seal due to facial hair, anatomy, blood or secretions/trauma; **O**bese or late pregnancy; **A**ge >55; **N**o teeth (roll gauze and place between gums and cheeks to improve seal); **S**tiff or increased airway pressures (asthma, COPD, obese, pregnant).

Difficult Laryngoscopy-LEMON: Look externally for anatomical distortions (small mandible, short neck, large tongue); Evaluate 3-3-2 Rule (Mouth open should accommodate 3 patient fingers, mandible to neck junction should accommodate 3 patient fingers, chin-neck junction to thyroid prominence should accommodate 2 patient fingers); **M**allampati (difficult to assess in the field); **O**bstruction / **O**bese or late pregnancy; **N**eck mobility.

Difficult Extraglottic Device Placement-RODS: Restricted mouth opening; **O**bstruction / **O**bese or late pregnancy; **D**istorted or disrupted airway; **S**tiff or increased airway pressures (asthma, COPD, obese, pregnant).

Nasotracheal intubation: *Orotracheal intubation is the preferred choice.* Procedure requires patient to have spontaneous breathing. Contraindicated in anatomically disrupted or distorted airways, increased intracranial pressure, severe facial trauma, basal skull fracture, head injury.


Pearls

- Consider preoxygenation/lung denitrogenation with a non-rebreather, a nasal cannula at 15 LPM, or CPAP prior to intubation (as patient condition allows).
- Severe hypotension (SBP<90) should be addressed with IV fluids and/or pressors (as appropriate) prior to intubation in order to reduce the likelihood of post-intubation cardiovascular decline.
- Capnometry (Color) or capnography is mandatory with all methods of intubation. Document results.
- Continuous capnography (ETCO₂) is mandatory for the monitoring of all patients with an ET tube.
- If an effective airway is being maintained by BVM and/or basic airway adjuncts (e.g. nasopharyngeal airway) with continuous pulse oximetry values of ≥90% or values expected based on pathophysiologic condition with otherwise reassuring vital signs (e.g. pulse oximetry of 85% with otherwise normal vitals in a post-drowning patient), it is acceptable to continue with basic airway measures instead of using an extraglottic airway device or intubation. Consider CPAP as indicated by protocol and patient condition.
- For the purposes of this protocol, a secure airway is achieved when the patient is receiving appropriate oxygenation and ventilation.
- An intubation attempt is defined as passing the laryngoscope blade or endotracheal tube past the teeth or inserted into the nasal passage.
- An appropriate ventilatory rate is one that maintains an ETCO₂ of 35 - 45. Avoid hyperventilation.
- Paramedics should use an extraglottic airway device if oral-tracheal intubation is unsuccessful.
- Maintain C-spine stabilization for patients with suspected spinal injury.
- Cricoid pressure and BURP maneuver may assist with difficult intubations. They may worsen view in some cases.
- Gastric tube placement should be considered in all intubated patients, if time allows.
- It is important to secure the endotracheal tube well.

PEDIATRIC TREATMENT PROTOCOLS

(for patients under 12 years of age)

General Pediatric Assessment



Pearls

- For all scenes where patient needs exceed available EMS resources, initial assessment and treatment shall be in accordance with an approved triage methodology.
- Correct life-threatening problems as identified.
- If the ability to adequately ventilate a patient cannot be established, the patient must be transported to the nearest emergency department.
- Never withhold oxygen from a patient in respiratory distress.
- Contact with online medical control should be established by radio. Telephone contact may only be used if the call is routed via a recorded phone patch through the FAO at 702-382-9007.


Disposition

- Patients sustaining traumatic injuries shall be transported in accordance with the Trauma Field Triage Criteria Protocol.
- Patients sustaining burn injuries shall be transported in accordance with the Burns Protocol.
- Pediatric patients (<18 y/o for transport purposes only) shall be transported in accordance with the Pediatric Destination Protocol.
- Patients with evidence of a stroke shall be transported in accordance with the Stroke Protocol.
- Sexual assault victims <13 y/o shall be transported to Sunrise Hospital.
- Sexual assault victims 13 y/o up to 18 y/o shall be transported to Sunrise Hospital or UMC.
- Sexual assault victims 18 y/o and older shall be transported to UMC.
- For sexual assault victims outside a 50-mile radius from the above facilities, the patient shall be transported to the nearest appropriate facility.
- Stable patients shall be transported to the hospital of their choice; if the patient has no preference, the patient should be transported to the closest facility.

Internal Disaster

- If a hospital declares an internal disaster, that facility is to be bypassed for all patients except patients in cardiac arrest, or in whom the ability to adequately ventilate has not been established.
- Operational exceptions may be initiated in regard to transport to hospitals on internal disaster.

General Pediatric Trauma Assessment



History

- Time and mechanism of injury
- Damage to structure or vehicle
- Location in structure or vehicle
- Others injured or dead
- Speed and details of MVC
- Restraints/protective equipment
- Past medical history
- Medications

Signs and Symptoms

- Pain, Swelling
- Deformity, lesions, bleeding
- AMS or unconscious
- Hypotension or shock
- Arrest


Differential (Life threatening)

- Tension pneumothorax
- Flail chest
- Pericardial tamponade
- Open chest wound
- Hemothorax
- Intra-abdominal bleeding
- Pelvis/femur fracture
- Spine fracture/cord injury
- Head injury
- Extremity fracture
- HEENT (airway obstruction)
- Hypothermia

Pearls

- Recommended exam: Mental Status, Skin, HEENT, Heart Lung, Abdomen, Extremities, Back, Neuro.
- Transport destination is based on the Trauma Field Triage Criteria Protocol.
- Transport should not be delayed for procedures; ideally procedures should be performed enroute when possible.
- BVM is an acceptable method of ventilating and managing an airway if pulse oximetry can be maintained $\geq 90\%$.
- Pediatric patients should be evaluated with a high index of suspicion; occult injuries may be present and pediatric patients can decompensate quickly.

Pediatric Abdominal Pain, Nausea & Vomiting



****Round up to nearest ½ pill**

History

- Age
- Medical/surgical history
- Onset
- Quality
- Severity
- Fever

Signs and Symptoms

- Pain location
- Tenderness
- Nausea
- Vomiting
- Diarrhea
- Dysuria
- Constipation


Differential

- Liver (Hepatitis)
- Gastritis
- Pancreatitis
- Kidney stone
- Appendicitis
- Bladder
- Bowel obstruction
- Gastroenteritis

Pearls

- Recommended Exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Back, Extremities, Neuro.
- Document mental status and vital signs prior to administration of antiemetics & pain management.
- Repeat vital signs after each fluid bolus.
- Consider retroperitoneal palpation for kidney pain.
- Pediatric fluid bolus is 20 ml/kg; may repeat to a maximum of 60 ml/kg.
- If there is suspicion that the patient is in DKA, do not exceed 20 ml/kg NS or LR.
- Morphine is not recommended in children for abdominal pain.
- Consider cardiac and ETCO₂ monitoring.

Pediatric Allergic Reaction



History

- Onset and location
- Insect sting or bite
- Food allergy/exposure
- Medication allergy/exposure
- New clothing, soap, detergent
- Past history of reactions
- Past medical history
- Medication history

Signs and Symptoms

- Itching or hives
- Coughing/wheezing or respiratory distress
- Throat or chest constriction
- Difficulty swallowing
- Hypotension/shock
- Edema
- Nausea/vomiting

Differential

- Urticarial (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration/airway obstruction
- Asthma/COPD
- CHF

Pearls

- Recommended Exam: Mental Status, Skin, Heart, Lung.
- Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.
- Epinephrine is a first-line drug that should be administered in acute anaphylaxis (moderate/severe symptoms). IM Epinephrine (1:1,000) should be administered in priority before or during attempts at IV or IO access.
- Contact Medical Control for refractory anaphylaxis.

Severity

- **Mild** reactions involve skin rashes, itchy sensation or hives with no respiratory involvement.
- **Moderate** reactions involve skin disorders and may include some respiratory involvement like wheezing, yet the patient still maintains good tidal volume air exchange.
- **Severe** reactions involve skin disorders, respiratory difficulty, and may include hypotension.


Special Considerations

- Always perform ECG monitoring when administering Epinephrine.
- Consider Dopamine for hypotension refractory to administration of Epinephrine.
- Provide oxygen and airway support as needed.

QI Metrics

- Epinephrine given appropriately.
- Airway assessment documented.

Pediatric Altered Mental Status



History

- Past medical history
- Medications
- Recent illness
- Irritability
- Lethargy
- Changes in feeding/sleeping
- Diabetes
- Potential ingestion
- Trauma

Signs and Symptoms

- Decrease in mentation
- Change in baseline mentation
- Decrease in blood sugar
- Cool, diaphoretic skin
- Increase in blood sugar
- Warm, dry, skin; fruity breath
- Kussmaul respirations, signs of dehydration

Differential


- Hypoxia
- CNS (trauma, stroke, seizure, infection)
- Thyroid (hyper/hypo)
- Shock (septic-infection, metabolic, traumatic)
- Diabetes (hyper/hypoglycemia)
- Toxicological
- Acidosis/Alkalosis
- Environmental exposure
- Electrolyte abnormalities
- Psychiatric disorder

Pearls

- Recommended Exam: Mental Status, HEENT, Skin, Heart, Lung, Abdomen, Back, Extremities and Neuro.
- Pay careful attention to the head exam for signs of injury.
- Be aware of AMS as presenting sign of an environmental toxin or Haz-Mat exposure and protect personal safety and that of other responders.
- Consider alcohol, prescription drugs, illicit drugs and over the counter preparations as possible etiology.
- If narcotic overdose or hypoglycemia is suspected, administer Narcan 0.1 mg/kg or Glucose prior to advanced airway procedures.
- Narcan is not recommended in the newly born.

Pediatric Behavioral Emergency

Protocol for pediatric patients <12 yrs old



History

- Situational crisis
- Psychiatric illness/medications
- Injury to self or threat to others
- Medic Alert tag
- Substance abuse/overdose
- Diabetes

Signs and Symptoms

- Anxiety, agitation, confusion
- Affect change, hallucinations
- Delusional thoughts, bizarre behavior
- Combative, violent
- Expression of suicidal/homicidal thoughts

Differential

- Altered mental status differential
- Alcohol intoxication
- Toxin/substance abuse
- Medication effect or overdose
- Withdrawal syndromes
- Depression
- Bipolar
- Schizophrenia
- Anxiety disorder

Pearls

- Midazolam is **NOT** recommended for use in children for behavioral emergencies.
- Law enforcement assistance should be requested on all calls involving potentially violent patients.
- Under no circumstances are patients to be transported restrained in the prone position.
- Recommended Exam: Mental Status, Skin, Heart, Lung, Neuro.
- Consider all possible medical/trauma causes for behavior.
- Do not irritate the patient with a prolonged exam.
- EMS providers are mandatory reporters in regard to suspected abuse of any vulnerable person.
- Consider cardiac and ETCO₂ monitoring.


Dystonic Reaction

- Condition causing involuntary muscle movements or spasms typically of the face, neck and upper extremities.
- Typically an adverse reaction to drugs such as Haloperidol (may occur with administration).
- When recognized, administer Diphenhydramine 1 mg/kg up to 50 mg IM/IV.

S.A.F.E.R.

- Stabilize the situation by containing and lowering the stimuli.
- Assess and acknowledge the crisis.
- Facilitate the identification and activation of resources (chaplain, family, friends or police).
- Encourage patient to use resources and take actions in his/her best interest.
- Recovery or referral – leave patient in care of responsible person or professional, or transport to appropriate facility.

Pediatric Bradycardia



History

- Respiratory insufficiency
- Past medical history
- Medications
- Pacemaker

Signs and Symptoms

- HR <60/min with hypotension, acute altered mental status, chest pain, acute CHF, seizures, syncope, or shock secondary to bradycardia
- Respiratory distress

Differential

- Hypoxia
- Hypothermia
- Sinus bradycardia
- Athletic
- Head injury (elevated ICP)
- Spinal cord lesion
- Overdose

Pearls

- Pediatric pacing is by Telemetry Physician order only.
- Recommended Exam: Mental Status, HEENT, Heart, Lung, Neuro.
- Bradycardia causing symptoms is typically <50/minute. Rhythm should be interpreted in the context of symptoms and pharmacological treatment given only when symptomatic; otherwise, monitor and reassess.
- Identifying signs and symptoms of poor perfusion caused by bradycardia are paramount.
- Hypoxemia is a common cause of bradycardia; be sure to oxygenate the patient and provide ventilatory support as needed.

Pediatric Burns



History

- Type of exposure (heat, gas, chemical)
- Inhalational injury
- Time of injury
- Past medical history & medications
- Other trauma
- Loss of consciousness
- Tetanus/immunization status

Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/distress
- Wheezing
- Singed facial or nasal hair
- Hoarseness or voice changes

Differential

- Superficial (1st degree) – red and painful
- Partial Thickness (2nd degree) – blistering
- Full Thickness (3rd degree) – painless/charred or leathery skin
- Thermal
- Chemical
- Electrical
- Radiation
- Lightning

Pearls

- Burn patients are trauma patients; evaluate for multisystem trauma.
- Assure whatever has caused the burn, is no longer contacting the injury. (Stop the burning process!)
- Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, Neuro.
- Consider early intubation with patients experiencing significant inhalation injuries.
- Potential CO exposure should be treated with 100% oxygen. (For patients in which the primary event is CO inhalation, transport to a hospital equipped with a hyperbaric chamber is indicated [when reasonably accessible].)
- Circumferential burns to extremities are dangerous due to potential vascular compromise secondary to soft tissue swelling. Elevate extremity.
- Burn patients are prone to hypothermia - never apply ice or cool burns; must maintain normal body temperature.
- Consider ETCO₂ monitoring.
- Evaluate the possibility of child abuse with children and burn injuries

Patients meeting the following Criteria shall be transported to the closest appropriate Burn Care Center:


1. Second degree burns >10% body surface area (BSA).
2. Any Third degree burns.
3. Burns that involve the face, hands, feet, genitalia, perineum, or major joints.
4. Electrical burns including lightning injury.
5. Chemical burns.
6. Circumferential burns.
7. Inhalation burns.
8. Burn injury with concomitant trauma

Pearls (Electrical)

- Do not contact the patient until you are certain the source of the electric shock has been disconnected.
- Attempt to locate contact points, (entry wound where the AC source contacted the patient; an exit at the ground point); both sites will generally be full thickness.
- Cardiac monitor; anticipate ventricular or atrial irregularity to include V-Tach, V-Fib, heart blocks, etc.
- Attempt to identify the nature of the electrical source (AC vs DC), the amount of voltage and the amperage the patient may have been exposed to during the electrical shock.

Pearls (Chemical)

- Certainly 0.9% NaCl Sol'n or Sterile Water is preferred; however if it is not readily available, do not delay; use tap water for flushing the affected area or other immediate water sources. Flush the area as soon as possible with the cleanest, readily available water or saline solution using copious amounts of fluids.



Early Intubation Indications

- Signs of Airway Obstruction
- Hoarseness, Stridor, Dysphagia
- Extensive Deep Facial Burns
- Signs of Respiratory Compromise
 - Accessory Muscle Use
 - Inability to Clear Secretions
 - Poor Oxygenation
- Burns in Mouth
- Total BSA ≥ 40%
- Altered Mentation
- Significant Risk of Edema

Fluid Resuscitation

- 13 years and above 500 ml NS or LR bolus
- 6-12 years 250 ml NS or LR bolus
- 5 years or less 125 ml NS or LR bolus
- Contact Burn Center Medical Direction for additional boluses or drip rates or if it is a prolonged transport.

History

- Events leading to arrest
- Estimated down time
- Past medical history
- Medications
- Existence of terminal illness

Signs and Symptoms

- Unresponsive
- Apneic
- Pulseless

Differential

- Medical vs. Trauma
- VF vs. Pulseless VT
- Asystole
- PEA
- Primary cardiac event vs. respiratory or drug overdose


Pearls

- Respiratory failure resulting in cardiac arrest should be addressed as it is identified.
- Efforts should be directed at high quality chest compressions with limited interruptions and early defibrillation when indicated.
- Consider early IO placement if IV is difficult.
- DO NOT HYPERVENTILATE.
- Reassess and document ETT placement using auscultation and ETCO₂ capnography.
- Switch compressors every two minutes.
- Try to maintain patient modesty.
- Mechanical chest compression devices should be used if available and appropriate for patient age/size in order to provide for consistent uninterrupted chest compressions and crew safety.
- Adult paddles/pads may be used on children weighing greater than 10 kg.
- Pre-assignment of pit crew roles is recommended. When this is not possible, tasks may be assigned by order of arrival: 1st: Airway; 2nd: Compressions; 3rd: IV/IO access, medication administration; 4th: Measure, Monitor/AED placement; 5th: family liaison/history gathering
- Pre-plan drug dosing based on weight estimations while en route and verify with a height based tape once reaching the patient
- Proper BVM selection: <5 kg = infant BVM. 5-30 kg = pediatric BVM. >30 kg = adult BVM.

H's & T's (reversible causes)

- Hypovolemia – Volume infusion
- Hypoxia – Oxygenation & ventilation, CPR
- Hydrogen ion (acidosis) – Ventilation, CPR
- Hyperkalemia – Calcium Chloride, Glucose, Sodium Bicarbonate, Albuterol
- Hypokalemia
- Hypothermia – Warming
- Hypoglycemia – Glucose
- Tension pneumothorax – Needle decompression
- Tamponade, cardiac – Volume infusion
- Toxins – Agent specific antidote
- Thrombosis, pulmonary – Volume infusion
- Thrombosis, coronary – Emergent PCI

Pediatric Cold-Related Illness



History

- Age, very young and old
- Exposure to decreased temperatures, but may occur in normal temperatures
- Past medical history/medications
- Drug or alcohol use
- Infections/sepsis
- Time of exposure/wetness/wind chill

Signs and Symptoms

- AMS/coma
- Cold, clammy
- Shivering
- Extremity pain
- Bradycardia
- Hypotension or shock

Differential

- Sepsis
- Environmental exposure
- Hypoglycemia
- Stroke
- Head injury
- Spinal cord injury

Pearls

- Recommended exam: Mental Status, Heart, Lung, Abdomen, Extremities, Neuro.
- Extremes of age are more prone to cold emergencies.
- Obtain and document patient temperature.
- If temperature is unknown, treat the patient based on suspected temperature.
- Hot packs can be used on the armpit and groin; care should be taken not to place the packs directly on the skin.

Hypothermia Categories

- Mild 90° - 95° F (33° - 35° C)
- Moderate 82° - 90° F (28° - 32° C)
- Severe <82° F (<28° C)


Hypothermia Mechanisms

- Radiation
- Convection
- Conduction
- Evaporation

Active Heating Measures

- Hot packs to the armpits and groin (do not place directly onto the skin)

Pediatric Drowning



History

- Submersion in liquid regardless of depth
- Possible history of trauma (dive)
- Duration of immersion
- Temperature of water or possibility of hypothermia
- Degree of water contamination

Signs and Symptoms

- Unresponsive
- Mental status changes
- Decreased or absent vital signs
- Vomiting
- Coughing, wheezing, rales, stridor, rhonchi
- Apnea
- Frothy/foamy sputum

Differential

- Trauma
- Pre-existing medical condition
- Barotrauma
- Decompression illness
- Post-immersion syndrome


Pearls

- Recommended Exam: Trauma Survey, Head, Neck, Chest, Abdomen, Back, Extremities, Skin, Neuro.
- Ensure scene safety.
- Hypothermia is often associated with submersion incidents.
- All patients should be transported for evaluation because of potential for worsening over the next several hours.

QI Metrics

- Complete and submit the SNHD Submersion Incident Report Form.

Pediatric Epistaxis



History

- Age
- Past Medical History
- Medications (HTN, Anticoagulants, aspirin, NSAIDS)
- Previous episodes of epistaxis
- Trauma
- Duration of bleeding
- Quantity of bleeding

Signs and Symptoms

- Bleeding from nasal passages
- Pain
- Nausea
- Vomiting

Differential


- Trauma
- Infection (viral URI or Sinusitis)
- Allergic rhinitis
- Lesions (polyps, ulcers)
- Hypertension

Oxymetazoline (Afrin) and Phenylephrine (Neosynephrine) should be avoided if child cannot follow instructions to blow their nose, or are unable to tolerate the administration of a nasal medication.

Pearls

- Recommended exam: Mental Status, HEENT, Lungs, Neuro
- History should include any clotting disorders such as Hemophilia or Von Willebrand disease, as these can contribute to bleeding.
- Non-accidental trauma as well as foreign body should be considered in pediatric patients with epistaxis.
- It is very difficult to quantify the amount of blood loss with epistaxis
- Bleeding may be also occurring posteriorly. Evaluate for posterior blood loss by examining the posterior pharynx.
- Detailed medication history should be obtained to assess for any NSAIDS, Antiplatelet agents or Anticoagulant medications that may contribute to bleeding.

Pediatric Heat-Related Illness



History

- Age, very old and young
- Exposures to increased temperatures and/or humidity
- Past medical history/medications
- Time and duration of exposure
- Poor PO intake, extreme exertion
- Fatigue and/or muscle cramping

Signs and Symptoms

- AMS/coma
- Hot, dry, or sweaty skin
- Hypotension or shock
- Seizures
- Nausea

Differential

- Fever
- Dehydration
- Medications
- Hyperthyroidism
- DTs
- Heat cramps, heat exhaustion, heat stroke
- CNS lesions or tumors

Pearls

- Recommended exam: Mental Status, Skin, Heart, Lung, Abdomen, Extremities, Neuro.
- Extremes of age are more prone to heat emergencies.
- Cocaine, amphetamines, and salicylates may elevate body temperatures.
- Sweating generally disappears as body temperatures rise over 104° F (40° C).
- Intense shivering may occur as patient is cooled.
- Active cooling includes application of cold packs or ice (not directly on skin), fanning either by air conditioning or fanning.
- Cold Saline is not to be administered for the treatment of hyperthermia unless directed by telemetry physician.

Heat Cramps

- Consist of benign muscle cramping caused by dehydration and is not associated with an elevated temperature.

Heat Exhaustion

- Consists of dehydration, salt depletion, dizziness, fever, AMS, headache, cramping, N/V. Vital signs usually consist of tachycardia, hypotension and elevated temperature.


Heat Stroke

- Consists of dehydration, tachycardia, hypotension, temperature >104° F (40° C) and AMS.

Active Cooling Measures

- Cold packs
- Ice (do not place directly on the skin)
- Fanning
- Air conditioning

Neonatal Resuscitation



Caveats:

- Deep airway suctioning is no longer recommended
- Traditional CPR 3:1 ratio is standard for newborns
- Most newborns requiring resuscitation will respond to BVM, compressions and Epi; for those that don't, consider hypovolemia, pneumothorax, and/or hypoglycemia (BG <40)

History

- Due date
- Time contractions started/ duration/frequency
- Rupture of membranes (meconium)
- Time and amount of any vaginal bleeding
- Sensation of fetal movement
- Prenatal care
- Past medical and delivery history
- Medications
- Gravida/Para Status
- High risk pregnancy

Signs and Symptoms

- Spasmodic pain
- Vaginal discharge or bleeding
- Crowning or urge to push
- Meconium

Differential

- Abnormal presentation (breech, limb)
- Prolapsed cord
- Placenta previa
- Abruptio placenta

Pearls


- Recommended Exam: Mental Status, Skin, HEENT, Neck, Chest, Heart, Lungs, Abdomen, Neuro.
- Document all times (delivery, contraction, duration, frequency).
- Some bleeding is normal; copious amounts of blood or free bleeding is abnormal.
- Record APGAR at one and five minutes after birth.
- APGAR of 7-10 is normal, while 4-7 require resuscitative measures
- Transport mother and infant together whenever possible.

APGAR	Score=0	Score=1	Score=2
• Activity/Muscle Tone	Absent	Arms/legs flexed	Active Movement
• Pulse	Absent	Below 100	Above 100
• Grimace/Reflex Irritability	No response	Grimace	Sneeze, cough, pulls away
• Appearance/Skin Color	Blue-Grey, pale all over	Normal, except extremities	Normal over entire body
• Respiration	Absent	Slow, irregular	Good, crying

Caveats:

- Deep airway suctioning no longer recommended.
- Traditional CPR 3:1 ratio is standard for newborns.
- Most newborns requiring resuscitation will respond to BVM, compressions and Epinephrine; for those that don't, consider hypovolemia, pneumothorax, and/or hypoglycemia (BG <40).

Pediatric Overdose / Poisoning



History

- Ingestion or suspected ingestion of a potentially toxic agent
- Substance ingested, route, quantity
- Time of ingestion
- Reason (suicidal, accidental, criminal)
- Available medications in home
- Past medical history, medications

Signs and Symptoms

- Mental status changes
- Hypotension/hypertension
- Decreased respiratory rate
- Tachycardia, dysrhythmias
- Seizures
- SLUDGE
- Malaise, weakness
- GI symptoms
- Dizziness
- Syncope
- Chest pain

Differential

- TCA overdose
- Acetaminophen OD
- Aspirin
- Depressants
- Stimulants
- Anticholinergic
- Cardiac medications
- Solvents, alcohols, cleaning agents, insecticides


Pearls

- Pediatric patients should be evaluated by a physician if an overdose/poisoning is suspected regardless of agent, amount or time.
- Recommended exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Neuro.
- Calcium Chloride is contraindicated in patients taking digitalis products.
- Overdose or toxin patients with significant ingestion/exposure should be closely monitored and aggressively treated. Do not hesitate to contact medical control if needed.
- In the case of cyanide poisoning, altered mental status may be profound. Profound altered mental status can be defined as a deficit that includes disorientation, bewilderment and difficulty following commands.
- Poison Control: 1-800-222-1222

Agents

- Acetaminophen: Initially normal or N/V. Tachypnea and AMS may occur later. Renal dysfunction, liver failure and/or cerebral edema may manifest.
- Depressants: Decreased HR, BP, temp and RR.
- Anticholinergic: Increased HR, increased temp, dilated pupils and mental status changes.
- Insecticides: May include S/S of organophosphate poisoning.
- Solvents: N/V, cough, AMS.
- Stimulants: Increased HR, BP, temp, dilated pupils, seizures and possible violence.
- TCA: Decreased mental status, dysrhythmias, seizures, hypotension, coma, death.

Pediatric Pain Management



History

- Age
- Location, duration
- Severity (1-10)
- Past medical history
- Pregnancy status
- Drug allergies and medications

Signs and Symptoms

- Severity (pain scale)
- Quality
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

Differential

- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural, respiratory
- Neurogenic
- Renal (colic)


Pearls

- Recommended exam: Respiratory Status, Mental Status, Area of pain, Neuro.
- Pain severity (1-10) is a vital sign to be recorded before and after medication administration and patient hand off.
- Monitor BP and respirations closely as sedative and pain control agents may cause hypotension and or respiratory depression.
- Consider patient's age, weight, clinical condition, use of drugs/alcohol, exposure to opiates when determining initial opiate dosing. Weight based dosing may provide a standard means of dosing calculation but it does not predict response.
- Exercise caution when administering opiates and benzodiazepines; this combination results in deeper anesthesia with significant risk of respiratory compromise.
- Burn patients may require more aggressive dosing.

QI Metrics

- Vital signs with O₂ sats documented.
- Pain scale documented before and after each intervention.
- Repeat vital signs after each intervention.

Pediatric Respiratory Distress



History

- Asthma
- Home treatment (oxygen, nebulizers)
- Medication
- Toxic exposure

Signs and Symptoms

- Shortness of breath
- Pursed lip breathing
- Decreased ability to speak
- Increased respiratory rate and effort
- Wheezing, rhonchi
- Use of accessory muscles
- Fever, cough
- Tachycardia


Differential

- Asthma
- Anaphylaxis
- Aspiration
- Pleural effusion
- Pneumonia
- Pneumothorax
- Pericardial tamponade (trauma)
- Hyperventilation
- Inhaled toxin

Pearls

- Be prepared to assist ventilations as needed.
- Recommended exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro.
- Pulse oximetry and end tidal continuous waveform capnography must be monitored.
- Consider MI.
- Allow the patient to assume a position of comfort.

Pediatric Seizure



History

- Reported or witnessed seizure activity
- Previous seizure history
- Seizure medications
- History of trauma
- History of diabetes
- Time of seizure onset
- Number of seizures
- Alcohol use, abuse or abrupt cessation
- Fever

Signs and Symptoms

- Decreased mental status
- Sleepiness
- Incontinence
- Observed seizure activity
- Evidence of trauma
- Unconsciousness

Differential

- CNS trauma
- Tumor
- Metabolic, hepatic or renal failure
- Hypoxia
- Electrolyte abnormality (Na, Ca, Mg)
- Drugs, medications non-compliance
- Infection, fever
- Alcohol withdrawal
- Hyperthermia
- Hypothermia

Pearls

- Recommended exam: Mental Status, HEENT, Heart, Lungs, Extremities, Neuro.
- Benzodiazepines are effective in terminating seizures; do not delay IM/IN administration while initiating an IV.
- Status epilepticus is defined as two or more seizures successively without an intervening lucid period, or a seizure lasting over five minutes.
- Grand mal seizures (generalized) are associated with loss of consciousness, incontinence and oral trauma.
- Focal seizures affect only part of the body and are not usually associated with a loss of consciousness.
- Be prepared to address airway issues and support ventilations as needed.
- Consider cardiac and ETCO₂ monitoring.

Pediatric Shock

General Pediatric Assessment

- E** Oxygen 15 L NRB
- A** Vascular Access
- P** Cardiac monitor/Capnography

Alternative appropriate treatment protocols as indicated

Estimated Minimum Systolic BP Calculation
 $(\text{Age in Years} \times 2) + 70$

BP calculation applies up to age 10 years


Trauma - related

General Trauma

Non-Trauma related

A NS or LR bolus 20 ml/kg IV/IO; may repeat x 2 with no rales on lung exam

For patients with known adrenal insufficiency, administer patient's own Solu-Cortef (hydrocortisone) as prescribed



Continue General Pediatric Assessment

History

- Blood loss-vaginal bleeding, ectopic, GI bleeding or AAA
- Fluid loss-vomiting, diarrhea, fever
- Infection
- Cardiac tamponade
- Medications
- Allergic reaction
- Pregnancy
- History of poor oral intake

Signs and Symptoms

- Restlessness, confusion
- Weakness, dizziness
- Weak rapid pulse
- Pale, cool, clammy skin
- Delayed capillary refill
- Hypotension
- Coffee-ground emesis
- Tarry stools

Differential

- Hypovolemic shock
- Cardiogenic shock
- Septic shock
- Neurogenic shock
- Anaphylactic shock
- Ectopic pregnancy
- Dysrhythmias
- Pulmonary embolism
- Tension pneumothorax
- Medication effect or overdose
- Vasovagal

For patients with known adrenal insufficiency, administer patient's own Solu-Cortef (hydrocortisone) as prescribed.

Causes of Adrenal Insufficiency:**Addison's Disease****Congenital Adrenal Hyperplasia****Long term administration of steroids****Others****Pearls**

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Hypotension can be defined as a systolic <Estimated Minimum Systolic. This is not always reliable and should be interpreted in context and patient's typical BP, if known. Shock may present with a normal BP initially.
- Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the only manifestation.
- Consider all possible causes of shock and treat per appropriate protocol.

Hypovolemic shock

- Hemorrhage, trauma, GI bleeding, ruptured aortic aneurysm or pregnancy-related bleeding

Cardiogenic shock

- Heart failure, MI, cardiomyopathy, myocardial contusion, toxins


Distributive shock

- Sepsis, anaphylaxis, neurogenic, toxins

Obstructive shock

- Pericardial tamponade, pulmonary embolus, tension pneumothorax

Pediatric Smoke Inhalation



History

- Exposed to smoke in a structure fire
- Exposed to smoke in a vehicle fire
- Exposed to smoke from other sources, industrial, confined space, wilderness fire etc.

Signs and Symptoms

- Facial burns
- Singed nasal hairs or facial hair
- Shortness of breath
- Facial edema
- Stridor
- Grunting respirations

Differential

- COPD
- CHF
- Toxic inhalation injury
- Caustic inhalation injury

Pearls

- Protect yourself and your crew.
- Have a high index of suspicion when treating patients at the scene of a fire.
- If the medication is not available on scene do not delay transport waiting for it.
- Carefully monitor respiratory effort and correct life threats immediately.
- Decide early on if you want to intubate as burned airways swell making intubation difficult.
- Profound altered mental status can be defined as a deficit that includes disorientation, bewilderment and difficulty following commands.

Pediatric Cyanokit Instructions and Dosing


1. Reconstitute Cyanokit vial per the Instructions. **The new vial concentration will be 25mg/ml.**
2. See chart below to find the appropriate reconstituted dose.

APPROXIMATE AGE	NB-1 m	2 m	4-6 m	8-10 m	1-1.5 y	2-2.5 y	3-3.5 y	4-5 y	5.5-7 y	7.5-8 y	8.5-10 y
WEIGHT IN LBS	7-9 lb	11 lb	13-15 lb	18-20 lb	22-24 lb	26-31 lb	33-40 lb	42-48 lb	53-62 lb	66-70 lb	71-80 lb
WEIGHT IN KGS	3-4 kg	5 kg	6-7 kg	8-9 kg	10-11 kg	12-14 kg	15-18 kg	19-22 kg	24-28 kg	30-32kg	34-36 kg
RECONSTITUTED SYRINGE AMOUNT	10 ml	15 ml	20 ml	25 ml	30 ml	40 ml	50 ml	60 ml	75 ml	85 ml	100 ml
TOTAL DOSAGE OF MEDICATION	250mg	375mg	500mg	625mg	750mg	1000mg	1250mg	1500mg	1875mg	2125mg	2500mg
gtts / sec	1	1	1	5	5	5	5	5	5	6	6

GIVE DOSE IN 50ml NS BAG **GIVE DOSE IN 250ml NS BAG**

3. Withdraw and waste the equivalent volume of Normal Saline from the bag size indicated.
4. Draw the appropriate dose from the vial referencing the chart, using the appropriate syringe size.
5. Inject the reconstituted medication into the appropriate sized bag of Normal Saline, below the Chart.
6. Spike the bag with 15 gtts/ml IV tubing.
7. Piggyback line into an IV/IO line and **Infuse over ~15 minutes** using the gtts/second noted above.

Pediatric Tachycardia / Stable (Normal Mental Status, Palpable Radial Pulse)



History

- Medications
- Diet (caffeine)
- Drugs (cocaine, methamphetamines)
- Past medical history
- Syncope/near syncope
- History of palpitations/racing heart

Signs and Symptoms

- Heart rate ≥ 180 in children
- Heart rate ≥ 220 in infants
- Dizziness, CP, SOB
- Diaphoresis


Differential

- Heart disease (WPW, valvular)
- Sick sinus syndrome
- Electrolyte imbalance
- Exertion, fever, pain, emotional stress
- Hypoxia
- Hypovolemia
- Drug effect, overdose
- Hyperthyroidism

Pearls

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Carefully monitor patients as you treat them; stable tachycardias may convert to unstable rhythms/conditions quickly.
- Sedate patients prior to cardioversion, if time allows.
- The most common tachyarrhythmia in children is sinus.

Pediatric Tachycardia / Unstable (Mental Status Changes, No Palpable Radial Pulse)



History

- Medications
- Diet (caffeine)
- Drugs (cocaine, methamphetamines)
- Past medical history
- Syncope/near syncope
- History of palpitations/racing heart

Signs and Symptoms

- Cardiac Arrest
- Heart rate ≥ 180 in children
- Heart rate ≥ 220 in infants
- Dizziness, CP, SOB
- Diaphoresis

Differential

- Heart disease (WPW, valvular)
- Sick sinus syndrome
- Electrolyte imbalance
- Exertion, fever, pain, emotional stress
- Hypoxia
- Hypovolemia
- Drug effect, overdose
- Hyperthyroidism

Pearls

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- If patient is in arrest, efforts should focus on quality chest compressions and rhythm correction.
- Administer Adenosine at a proximal IV site, rapidly followed by a saline flush.
- The most common tachyarrhythmia in children is sinus.

Pediatric Ventilation Management

Use supplemental oxygen to maintain an oxygen saturation of >94%; >90% for patients on home oxygen for chronic conditions

E	Basic Airway Maneuvers - Open airway, Chin lift/Jaw thrust - NPA or OPA as needed - Suction as needed	Respiratory Distress and/or Tracheostomy Tube Replacement Protocol if needed
	Consider Cervical Stabilization	
	Consider Altered Mental Status/Syncope	

E	Administer oxygen
	BVM as needed

Intervention effective? (Decision diamond)

A	Extraglottic Airway	Obtain Vascular Access
	Endotracheal Intubation	ECG Monitor
P	Consider Induction: Administer MIDAZOLAM 0.1 mg/kg IV/IN/IO Titrated to Effect. Maximum Single Dose : 5 mg. Must be Given Slowly Over 3-5 Minutes. Additional Doses by Physician Order Only. OR; KETAMINE 2 mg/kg IV/IO Or 4 mg/kg IM OR; ETOMIDATE 0.3mg/kg IV/IO. Max dose 20 mg	
	Post-Intubation Sedation Administer DIAZEPAM 0.2mg/kg IV/IO. Maximum dose 5 mg. May Repeat After 5 Minutes with Physician Order. Or; DIAZEPAM 0.5 mg/kg PR via #5 or #8 French feeding tube, Maximum Dose 20 mg	
	Post-Intubation Analgesia Administer FENTANYL 1 mcg/kg; max dose 100 mcg	

Extraglottic/ETT placement successful? (Decision diamond)

Able to ventilate without extraglottic/ETT in place? (Decision diamond)

P	Cricothyroidotomy
---	-------------------

Continue Care and Transport as Appropriate

Pearls

- Consider preoxygenation/lung denitrogenation with a non-rebreather or a nasal cannula at 15 LPM prior to intubation (as patient condition allows)
- Severe hypotension (SBP $70+(2 \times \text{age})$) should be addressed with IV fluids and/or pressors (as appropriate) prior to intubation in order to reduce likelihood of post-intubation cardiovascular decline.
- Capnometry (Color) or capnography is mandatory with all methods of intubation. Document results.
- Continuous capnography (ETCO₂) is mandatory for the monitoring of all patients with an ET tube.
- If an effective airway is being maintained by BVM and/or basic airway adjuncts (e.g. nasopharyngeal airway)
- with continuous pulse oximetry values of $\geq 90\%$, or values expected based on pathophysiologic condition
- with otherwise reassuring vital signs (e.g. pulse oximetry of 85% with otherwise normal vitals in a post drowning patient), it is acceptable to continue with basic airway measures instead of using an extraglottic airway device or intubation.
- For the purposes of this protocol, a secure airway is achieved when the patient is receiving appropriate oxygenation and ventilation.
- An Intubation Attempt is defined as passing the laryngoscope blade or endotracheal tube past the teeth or inserted into the nasal passage.
- An appropriate ventilatory rate is one that maintains an ETCO₂ of 35 - 45. Avoid hyperventilation.
- Paramedics should use an extraglottic airway device if oral-tracheal intubation is unsuccessful.
- Maintain C-spine stabilization for patients with suspected spinal injury.
- Cricoid pressure and BURP maneuver may assist with difficult intubations. They may worsen view in some cases.
- Gastric tube placement should be considered in all intubated patients if time allows.
- It is important to secure the endotracheal tube well.

OPERATIONS PROTOCOLS

Communications



Telemetry contact should be established by radio.
Telephone contact may only be used if the call is recorded via a
phone patch through the FAO at 702-382-9007.

1. Telemetry contact shall be established:
 - A. For all time sensitive or life threatening condition transports.
 - B. For any medical emergency in which the EMS provider's judgment suggests consultation with a telemetry physician is necessary.
 - C. For all trauma patients going to a trauma center.
 - D. When telemetry contact is required per protocol.
2. For patients who meet Trauma Field Triage Criteria, telemetry reports shall include:
 - A. ETA
 - B. Patient age
 - C. Gender
 - D. Mechanism of injury
 - E. Ambulatory at scene
 - F. Suspected injuries
 - G. Vital signs
 - H. Airway status
 - I. Neurologic status
 - J. An incident identifier if multiple patients are involved (e.g. fire department command code "Main Street Command")
3. Notify/meet with the receiving facility prior to transfer of care with suspected need for Contact Isolation Preparation
 - A. State the general type of agent involved (insect, chemical, biological, radiation, nuclear, explosive)
 - B. State the type of agent if known.
 - C. If unknown state the general type with patient symptoms. Example – "Unknown chemical substance causing respiratory distress with secretions.
4. For all other patients, telemetry reports shall include, at a minimum:
 - A. Attendant/vehicle identification
 - B. Nature of call: INFORMATION ONLY or REQUEST FOR PHYSICIAN ORDERS
 - C. Patient information (i.e. number, age, sex)
 - D. Patient condition (i.e. stable, full arrest)
 - E. History
 - 1) Basic problem or chief complaint
 - 2) Pertinent associated symptoms
 - 3) Time since onset
 - 4) Past history, if pertinent
 - F. Objective findings
 - 1) General status of patient
 - 2) Level of responsiveness
 - 3) Vital signs
 - 4) Pertinent localized findings
 - 5) Working impression of patient's problem
 - G. Treatment
 - 1) In progress
 - 2) Requests for drugs or procedures
 - H. Estimated time of arrival, including any special circumstances that may cause a delay in transport.
 - I. For patients meeting "Code White" or "Code STEMI" criteria, a preliminary telemetry report should be made to notify the receiving facility of the type of activation, and an estimated arrival time. An "Information Only" telemetry should follow once transport has been initiated.

Communications (Cont.)



Telemetry contact should be established by radio.
Telephone contact may only be used if the call is recorded via a
phone patch through the FAO at 702-382-9007.

5. Notification of transport shall be provided to the receiving hospital for ALL other calls.
 - A. Notification can be completed via:
 - 1) Radio
 - 2) Telephone
 - 3) EMSsystem
 - B. Notification reports shall include:
 - 1) Patient age
 - 2) Chief complaint
 - 3) Type of bed required (monitored/unmonitored)
 - 4) Unit number
 - 5) ETA
6. Providers will relay assessment findings and treatment provided to the individual(s) assuming responsibility for the patient(s).
7. Patient confidentiality must be maintained at all times.
8. All patients should be treated with dignity and respect in a calm and reassuring manner.

Do Not Resuscitate (DNR/POLST)

1. All patients with absent vital signs who do not have conclusive signs of death (refer to Prehospital Death Determination protocol) shall be treated with life-resuscitating measures unless EMS personnel are presented with a valid Do-Not-Resuscitate (DNR)/Physician Order for Life-Sustaining Treatment (POLST) Identification/Order.



- A. A valid DNR Identification is a form, wallet card, or medallion issued by the Southern Nevada Health District, Nevada Division of Public and Behavioral Health, or an identification issued by another state indicating a person's desire and qualification to have life-resuscitating treatment withheld.
- B. A valid DNR Order is a written directive issued by a physician licensed in this state that life-resuscitating treatment is not to be administered to a qualified patient. The term also includes a valid Do-Not-Resuscitate order issued under the laws of another state.
- C. A valid POLST form signed by a physician that records the wishes of the patient and directs a healthcare provider regarding the provision of life-resuscitating treatment and life-sustaining treatment.

Note: Verbal instructions from friends or family members *DO NOT* constitute a valid DNR/POLST.

2. In preparation for, or during a inter-facility transfer, a valid DNR Order/POLST in the qualified patient's medical record shall be honored in accordance with this protocol.
3. If the EMS provider is presented with a DNR/POLST Order or Identification, he shall attempt to verify the validity of the Order or Identification by confirming the patient's name, age, and condition of identification.
4. The DNR/POLST Order or Identification shall be determined *INVALID* if at any time the patient indicates that he/she wishes to receive life-resuscitating treatment. The EMS provider shall document the presence of the DNR/POLST Order or Identification, and how the patient indicated that he/she wanted the Order or Identification to be revoked. EMS personnel shall relay this information to any subsequent medical providers, including but not limited to, flight crews and staff at the receiving medical facility.
5. Once the DNR/POLST Order or Identification is determined to be valid and has not been revoked by the patient, the emergency care provider shall provide *ONLY* supportive care and withhold life-resuscitating treatment.
6. Faxed, copied or electronic versions of the DNR Identification/POLST are legal and valid.

Supportive Care:

- Suction the airway
- Administer oxygen
- Position for comfort
- Splint
- Control bleeding
- Provide pain medication (ALS only)
- Provide emotional support
- Contact hospice, home health agency, attending physician or hospital as appropriate
- Be attentive of any actions the patient may take to revoke his authorization to withhold life-resuscitating treatment

Do Not Resuscitate (DNR/POLST)(Cont.)

Withhold Life-Resuscitating Treatment:

CPR and its components including:

Chest compressions

Defibrillation

Cardioversion

Assisted ventilation

Airway intubation

Administration of cardiotoxic drugs

- 6. EMS personnel will document on the PCR the presence of the DNR/POLST Order or Identification. Documentation should include the patient's name, and the physician's name and identification number, which are found on the DNR/POLST Order or Identification.**
- 7. An EMS provider who is unwilling or unable to comply with the DNR protocol shall take all reasonable measures to transfer a patient with a DNR/POLST Order or Identification to another provider or facility in which the DNR/POLST protocol may be followed.**


Documentation

1. A Patient Care Record (PCR) will be completed for each incident/patient encounter, in accordance with current EMS Regulations. A patient is any individual that, upon contact with an EMS system, has any of the following:
 - A. A complaint or mechanism suggestive of potential illness or injury.
 - B. Obvious evidence of illness or injury.
 - C. An individual or informed 2nd/3rd party caller requests evaluation for potential illness or injury.

2. PCRs shall include no less than the following information:
 - A. Patient's name, address, age, and sex;
 - B. Date and location of call;
 - C. Time of dispatch, arrival at scene, departure from scene, and arrival at hospital;
 - D. Mechanism of injury—chief complaint;
 - E. Medication(s) used by patient and allergies;
 - F. Pertinent patient history, including current medication(s) and allergies;
 - G. Signs and symptoms identified during patient assessment, and changes;
 - H. Care and treatment given at scene and during transport;
 - I. Patient destination;
 - J. Name of attendant(s);
 - K. If care is provided as authorized by protocol;
 - L. File Attachments: The associated monitor file must be uploaded and attached to the PCR if the monitor was used for any of the following purposes:
 - 1) Assessing and/or monitoring the cardiac rhythm;
 - 2) Obtaining a 12-lead electrocardiogram (ECG)
 - 3) Providing electrical therapy; cardioversion, defibrillation, and/or pacing
 - 4) Monitoring End-Tidal Carbon Dioxide (ETCo2) levels and/or waveform of an intubated patient
 - M. In cases of trauma, the patient's trauma score, TFTC status, and any injury mitigation devices shall be documented, i.e. car seats, seat belts, air bags, helmets, etc.;
 - N. At least one full set of vital signs;
 - 1) Blood pressure
 - 2) Heart rate
 - 3) Respiratory rate
 - 4) Temperature as indicated
 - 5) Oxygen saturation
 - 6) Reassessment after interventions, i.e. pain score after medications;
 - 7) Any complications or other relevant information.

3. Any agency that provides patient care activities prior to the arrival of the transporting agency shall provide the transporting agency with, at a minimum, a verbal report reflecting those activities. This verbal report must be documented in the transporting agency's PCR.

Hostile Mass Casualty Incident



Inter-Facility Transport of Patients By Ambulance

1. Ambulance attendants should only transfer a patient whose therapy required during the transfer lies within the ambulance attendant's capabilities, unless capable personnel accompany the patient.
 - A. Ambulance attendants are authorized to administer or monitor all medications listed on the official drug inventory as appropriate for their level of licensure and as per protocol.
 - B. AEMT and Paramedic ambulance attendants are authorized to administer or monitor any crystalloid IV solution during transport.
 - C. EMT ambulance attendants are authorized to monitor locked intravenous peripheral lines during transport, including helpocks, Broviacs, Hickmans, Port-A-Catheters, and PICC lines.
 - D. Arterial lines should be discontinued unless appropriate personnel from the initiating facility accompany the patient.
 - E. Heparin locks/implantable catheters with/without reservoirs may be closed off and left in place. If they are to be used during transport, then an IV drip should be established if tolerated by the patient.
 - F. IV pump systems should be discontinued unless capable personnel accompany the patient.
 - G. Orogastric or nasogastric tubes may be left in place and should either be closed off or left to suction per order of the transferring physician.
 - H. Orthopedic devices may be left in place at the ambulance attendant's discretion as to ability to properly transport the patient with existing device(s) in place.
 - I. Trained personnel authorized to operate the apparatus should accompany any patient requiring mechanical ventilation during transport. If the patient will require manual ventilatory assistance, then at least two persons shall be available to attend to the patient.
2. Prior to the transfer, the transferring physician is responsible for notifying the receiving physician of the following:
 - 1) Reason for transfer
 - 2) Patient condition
 - 3) Estimated time of arrival
3. The transferring physician must provide the ambulance attendants with the name of the receiving facility and receiving physician, copies of any available diagnostic tests, X-rays, medical records, copy of code status, DNR, POLST, or advanced directive paperwork as applicable, any isolation precaution information, and the EMTALA form prior to releasing the patient.
4. Any agency that provides patient care activities prior to the arrival of the transporting agency shall provide the transporting agency with, at a minimum, a verbal report reflecting those activities. This verbal report must be documented in the transporting agency's PCR.

Pediatric Patient Destination

Pediatric patients (<18 years of age) shall be transported in accordance with the following criteria:

- 1. Pediatric patients (including psychiatric patients) shall be transported, based on the preference of the parent or legal guardian, to one of the following facilities:**
 - A. St. Rose Dominican – Siena Campus**
 - B. Summerlin Hospital Medical Center**
 - C. Sunrise Hospital & Medical Center**
 - D. University Medical Center**
- 2. If the parent or legal guardian does not have a preference, the patient shall be transported to the closest of the above facilities.**
- 3. If, in the judgment of prehospital personnel, the transport time to one of the above facilities would be detrimental to a critically ill/unstable pediatric patient, the patient should be transported to the closest Emergency Department.**
- 4. The patient may be transported to a non-designated facility:**
 - A. At the request of the parent or legal guardian, and if the child is deemed stable by the EMS provider; or**
 - B. The incident is greater than 50 miles from the closest pediatric facility; and**
 - C. The receiving facility and physician are contacted and agree to accept the patient.**
- 5. Pediatric sexual assault victims shall be transported as follows:**
 - A. Victims <13 years of age shall be transported to Sunrise Hospital and Medical Center.**
 - B. Victims 13 years of age and up to 18 years of age shall be transported to either Sunrise Hospital & Medical Center or University Medical Center.**
 - C. For sexual assault victims outside a 50-mile radius from the above facilities, the licensee providing emergency medical care shall transport the patient to the nearest appropriate facility.**

Prehospital Death Determination



For all emergency scenes where patient needs exceed available EMS resources, initial assessment and treatment shall be in accordance with the START/SMART triage methodology.

1. Patients who appear to have expired will not be resuscitated or transported by Clark County EMS personnel if any of the following obvious signs of death are present:



- A. Body decomposition
- B. Decapitation
- C. Transection of thorax (hemicorpectomy)
- D. Incineration
- E. For other traumatic injuries suspected to be incompatible with life, medical control must be contacted for medical direction.

If there are any extenuating circumstances regarding access to patient, contact medical control.

OR if ALL (5) presumptive signs of death AND AT LEAST one (1) conclusive sign of death are identified.

The (5) presumptive signs of death that **MUST** be present are:

- 1) Unresponsiveness
- 2) Apnea
- 3) Pulselessness
- 4) Fixed, dilated pupils
- 5) For Non-Traumatic Arrests, Asystole in at least 2 (two) leads or a "No Shock Advised" prompt from an AED



Conclusive signs of death include:

- 1) Dependent lividity
- 2) Rigor mortis

If any of the findings are different than those described above, clinical death is not confirmed, and resuscitative measures should be immediately initiated or continued.

2. Once it has been determined that the patient has expired and resuscitation will not be attempted:
- A) Immediately notify the appropriate authority;
 - B) **DO NOT** leave a body unattended. You may be excused once a responsible person (i.e. Coroner's investigator, police, security, or family member) is present;
 - C) **DO NOT** remove any property from the body or the scene for any purpose;
 - D) **NEVER** transport/move a body without permission from the Coroner's office except for assessment or its protection.



If the body is in the public view and cannot be isolated, screened, or blocked from view, and is creating an unsafe situation with citizens/family, the body can be covered with a clean, **STERILE BURN SHEET** obtained from the EMS vehicle.

Public Intoxication

1. A person who is suspected to be intoxicated and has no other emergent need should be transported to an approved alcohol and drug abuse facility rather than a hospital's emergency department *IF* the patient meets *ALL* of the following criteria:

A. Patient is able to stand with minimal assistance of one or two people.

B. Vitals as follows:

1. Blood Pressure: Systolic: 90 – 180
Diastolic: 60 – 100
2. Pulse Rate: 60 – 120
3. Respiratory Rate: 12 – 22
4. Blood Glucose between 60 – 250
5. Glasgow Coma Score \geq 14
6. SPO2 >94% or 90% if smoker
7. No acute medical complications
8. No signs of trauma
9. No suspected head injury
10. Approval of the physician or medical staff upon assessment prior to transport to an alternative facility. Contact with the facility needs to be routed via recorded phone patch through FAO at 702-382-9007

All of the above parameters must be met and the patient must be clinically stable.

2. If there is *ANY* doubt about whether the person is in need of emergency medical care, the person should be transported to a receiving emergency facility.

* Approved Public Intoxication
Facilities:

WestCare

Crossroads of Southern Nevada
(CRSN)

Quality Improvement Review

When EMS or hospital personnel wish to have an incident involving patient care reviewed within the Clark County EMS system, the following steps shall be taken:

1. The person requesting a review of an incident should contact the designated representative of the agency/hospital involved to initiate the process. If after gathering appropriate information and discussing the incident both parties are satisfied a problem does not exist, nothing further needs to be done.
2. If either party would like to pursue an investigation of the incident, the "Southern Nevada Health District EMS Incident Report" should be completed and a copy should be forwarded to the OEMSTS.
3. Upon receipt of the "Southern Nevada Health District EMS Incident Report" OEMSTS staff will review the case, gather information from the agencies/hospitals involved, and evaluate the need for further investigation. The agency/hospital may be asked to conduct an internal investigation, involving their medical director when appropriate, and provide a summary of their findings to the OEMSTS.
4. The personnel involved in the incident may be interviewed by the EMS medical director or his designee and their agency/hospital medical director to gather additional information.
5. Upon completion of the investigation, a report will be prepared and given to the agency/hospital representatives involved. Direct communication between the agency/hospital and complainant is recommended with a brief written summary of actions taken provided to the OEMSTS.
6. A quarterly aggregate summary of the incidents reviewed by the OEMSTS will be prepared and reported at the Quality Improvement Directors and Medical Advisory Board meetings.
7. All documentation and correspondence regarding this quality improvement activity; to monitor, review, evaluate, and report on the necessity, quality, and level of care provided a patient is confidential pursuant to NRS 49.117 – NRS 49.123, NRS 49.265, NRS 450B.810 and NRS 629.061.

Termination of Resuscitation

1. Licensed EMS personnel are not obligated to continue resuscitation efforts that have been started by other persons at the scene if the patient meets the criteria listed in the Prehospital Death Determination protocol. This includes telephone CPR initiated by Emergency Medical Dispatchers.
2. Resuscitation should be terminated/ not initiated without telemetry contact if a valid DNR/POLST or physician written order is provided.
3. Resuscitation started in the field may be discontinued only by a telemetry physician order when the following conditions have been met:
 - A. For medical arrest, contact closest hospital for telemetry physician order:

The patient remains in persistent asystole or agonal rhythm after twenty (20) minutes of appropriate resuscitation, to include:

 - 1) CPR
 - 2) Effective ventilation with 100% oxygenation
 - 3) Administration of appropriate ACLS medications, if available.
 - 4) Confirm no organized rhythm or PEA<40, or a “No Shock Advised” on AED
 - B. For traumatic arrest, contact TRAUMA CENTER based on catchment for telemetry physician order:
 - 1) Open airway with basic life support measures
 - 2) Provide CPR and effective ventilations with 100% oxygenation for two (2) minutes
 - 3) Perform bilateral needle thoracentesis if tension pneumothorax suspected
 - 4) Confirm no organized rhythm or PEA <40, or a “No Shock Advised” on AED
 - C. The patient develops, or is found to have one of the following conclusive signs of death at any point during the resuscitative effort:
 - 1) Lividity
 - 2) Rigor mortis
4. When resuscitation has been terminated in the field, all medical interventions shall be left in place.
5. If possible, do not leave a body unattended. Once a responsible person (i.e. Coroner’s investigator, police, security, or family member) is present at the scene, you may be excused.
6. **NEVER** transport/move a body without permission from the Coroner’s office, except for assessment or its protection.



If the body is in the public view and cannot be isolated, screened, or blocked from view, and is creating an unsafe situation with citizens/family, the body can be covered with a clean, **STERILE BURN SHEET** obtained from the EMS vehicle.

NOTES: In rural or wilderness situations, EMS providers must make every effort to contact medical control, but resuscitation may be terminated in the field without medical control when any of the following have occurred:

- A. There has been no return of pulse despite greater than 20 minutes of CPR and effective ventilation (consider extending in the case of hypothermia or drowning)
- B. Transport to an emergency department will take greater than 40 minutes (consider extending in the case of hypothermia or drowning)
- C. The EMS providers are exhausted and it is physically impossible to continue the resuscitation.

Transport Destinations

1. Medically stable patients should be transported to their hospital of choice if the destination is not significantly beyond the primary response area of the transporting agency. If the patient does not have a preference, the patient should be transported to the nearest appropriate facility.
2. Patients sustaining traumatic injuries shall be transported in accordance with the Trauma Field Triage Criteria Protocol.
3. Patients sustaining burn injuries shall be transported in accordance with the Burns Protocol.
4. Pediatric patients (<18 years of age for transport purposes *ONLY*) shall be transported in accordance with the Pediatric Destination Protocol.
5. Patients with evidence of an acute cerebrovascular accident shall be transported in accordance with the Stroke Protocol.
6. Sexual assault victims shall be transported as follows:
 - A. Victims <13 years of age shall be transported to Sunrise Hospital & Medical Center.
 - B. Victims 13 years of age and up to 18 years of age shall be transported to either Sunrise Hospital & Medical Center or University Medical Center.
 - C. Victims 18 years of age and older shall be transported to University Medical Center.
 - D. For sexual assault victims outside a 50-mile radius from the above facilities, the licensee providing emergency medical care shall transport the patient to the nearest appropriate facility.
7. All medical patients in cardiac arrest or in whom the ability to adequately ventilate cannot be established should be transported to the closest facility.
8. If a hospital declares an *Internal Disaster*, that facility is to be bypassed for *ALL* patients except patients in cardiac arrest, or in whom the ability to adequately ventilate has not been established.
9. For patients outside a 50 mile radius from protocol designated transport destinations, the licensee providing emergency medical care shall transport the patient to the nearest appropriate facility.

Remote Outpatient Emergency Department Alternate Destination Criteria

1. Patients who require a medical or psychiatric evaluation and do not have evidence of any potentially life-threatening illness or injury at the time of transport may be transported to a remote outpatient emergency department if;
2. The patient has normal vital signs, unless accepted by the remote outpatient emergency department, telemetry contact is made, and;
3. The patient does not meet any of the following exclusions criteria:
 - A. Violent or uncooperative patients
 - B. Obstetric patients > 20 weeks gestation
 - C. Any patient in need of time-critical intervention that can be provided only at a hospital-based emergency department. For example, but not limited to STEMI, Stroke, or ACS.
 - D. Any condition covered by another destination directive:

Trauma Field Triage Criteria	Normal Vital Signs:
Stroke Protocol	Heart Rate 60-100
Burns Protocol	Respiratory Rate 10-20
Pediatric Destination Protocol	Systolic BP 100-180
Sexual Assault Victims	Diastolic BP 60-110
Cardiac Arrest	Room air pulse oximetry >94%
	Alert and oriented X4
4. Alternate transportation and destination decisions should be consistent with medical necessity and with consideration for patient preference when the patient's condition allows.

Trauma Field Triage Criteria

A licensee providing emergency medical care to a patient at the scene of an injury shall use the following procedures to identify and care for patients with traumas:

1. Step 1 – Measure vital signs and level of consciousness. If the patient's:

- A. Glasgow Coma Scale is 13 or less;
- B. Systolic blood pressure is less than 90 mm Hg; or
- C. Respiratory rate is less than 10 or greater than 29 breaths per minute (less than 20 in infant aged less than 1 year), or is in need of ventilatory support

the adult patient *MUST* be transported to a Level 1 or 2 center for the treatment of trauma in accordance with the catchment area designated. The pediatric patient *MUST* be transported to a pediatric center for the treatment of trauma.

2. Step 2 – Assess anatomy of injury. If the patient has:

- A. Penetrating injuries to head, neck, torso, or extremities proximal to elbow or knee;
- B. Chest wall instability or deformity (e.g. flail chest);
- C. Two or more proximal long-bone fractures;
- D. Crushed, degloved, mangled, or pulseless extremity;
- E. Amputation proximal to wrist or ankle;
- F. Pelvis fractures;
- G. Open or depressed skull fractures; or
- H. Paralysis

the adult patient *MUST* be transported to a Level 1 or 2 center for the treatment of trauma in accordance with the catchment area designated. The pediatric patient *MUST* be transported to a pediatric center for the treatment of trauma.

3. Step 3 – Assess mechanism of injury and evidence of high-energy impact, which may include:

- A. Falls
 - 1) Adults: greater than 20 feet (one story is equal to 10 feet)
 - 2) Children: greater than 10 feet or two times the height of the child
- B. High-risk auto crash
 - 1) Motor vehicle was traveling at a speed of at least 40 miles per hour immediately before the collision occurred;
 - 2) Intrusion, including roof: greater than 12 inches occupant site; greater than 18 inches any site;
 - 3) Ejection (partial or complete) from automobile;
 - 4) Motor vehicle rolled over with unrestrained occupant(s);
 - 5) Death in same passenger compartment
- C. Motorcycle crash greater than 20 mph
- D. Auto vs pedestrian/bicyclist thrown, run over, or with significant (greater than 20 mph) impact

The patient *MUST* be transported to a Level 1, 2, or 3 center for the treatment of trauma in accordance with the catchment area designated. For patients who are injured outside a 50-mile radius from a trauma center, the licensee providing emergency medical care shall call and consider transport to the nearest receiving facility.

Trauma Field Triage Criteria (Cont.)

4. Step 4 – Assess special patients

A. Older adults

- 1) Risk of injury/death increases after age 55 years
- 2) SBP less than 110 mm Hg might represent shock after age 65 years
- 3) Low impact mechanisms (e.g. ground level falls) might result in severe injury

B. Children should be triaged preferentially to a trauma center.

C. Anticoagulants and bleeding disorders: Patients with head injury are at high risk for rapid deterioration.

D. Burns

- 1) Without other trauma mechanisms: transport in accordance with the Burns protocol
- 2) With trauma mechanism: follow appropriate catchment guidelines for trauma. Trauma patients with burns falling into St Rose Siena catchment area will be transported to Sunrise Hospital, and those falling in the Mike O'Callaghan catchment area will be transported to UMC Hospital.

E. Pregnancy greater than 20 weeks

F. EMS provider judgment

The patient *MUST* be transported to a Level 1, 2, or 3 center for the treatment of trauma in accordance with the catchment area designated. For patients who are injured outside a 50-mile radius from a trauma center, the licensee providing emergency medical care shall call and consider transport to the nearest receiving facility.

The person licensed to provide emergency medical care at the scene of an injury shall transport a patient to a designated center for the treatment of trauma based on the following guidelines:

St. Rose Dominican Hospital - Siena Campus (Level 3 Trauma Center) Catchment Area

All trauma calls that meet Step 3 or in the provider's judgment meet Step 4 of the Trauma Field Triage Criteria Protocol or pediatric Step 4 and occur within the City of Henderson or the geographical area bordered by Interstate 15 to the west and Sunset road to the north, and the county line to the east, are to be transported to St. Rose Hospital – Siena Campus and the medical directions for the treatment of the patient must originate at that center;

Mike O'Callaghan Military Medical Center (Level 3 Trauma Center) Catchment Area

All trauma calls that meet Step 3 or in the provider's judgment meet Step 4 of the Trauma Field Triage Criteria Protocol or pediatric Step 4 and occur within the geographical area bordered by Pecos Road to the west, Interstate 15 to the west/northwest and Lake Mead Blvd to the south, and the county line to the east, are to be transported to Mike O' Callaghan Military Medical Center and the medical directions for the treatment of the patient must originate at that center;

Sunrise Hospital & Medical Center (Level 2 Trauma Center) Catchment Area

All adult trauma calls and pediatric Step 3 or 4 trauma calls that meet the Trauma Field Triage Criteria Protocol and occur within the geographical area bordered by Paradise Road to the west, Sahara Avenue to the north, Sunset Road to the south, and the county line to the east, are to be transported to Sunrise Hospital & Medical Center and the medical directions for the treatment of the patient must originate at that center;

In addition, adult trauma calls that meet Step 1 or 2 of the Trauma Field Triage Criteria Protocol and occur within the St. Rose Dominican Hospital – Siena Campus Catchment Area, City of Henderson, or the geographical area bordered by Paradise Road to the west continuing along that portion where it becomes Maryland Parkway, Sunset Road to the north, and the county line to the east, are to be transported to Sunrise Hospital & Medical Center and the medical directions for the treatment of the patient must originate at that center.

Trauma Field Triage Criteria (Cont.)

University Medical Center (Level 1 Trauma Center and Pediatric Level 2 Trauma Center) Catchment Area

All trauma calls that meet the Trauma Field Triage Criteria and occur within any other area of Clark County are to be transported to University Medical Center/Trauma and the medical directions for the treatment of the patient must originate at that center.

All pediatric Step 1 and Step 2 trauma calls that occur within Clark County are to be transported to University Medical Center/Trauma and medical directions for the treatment of the patient must originate at that center.

In addition, adult trauma calls that meet Step 1 or 2 of the Trauma Field Triage Criteria Protocol and occur in the geographical area bordered by Paradise road to the east, Sunset Road to the north, Interstate 15 to the west, and the county line to the south, are to be transported to University Medical Center/Trauma and the medical directions for the treatment of the patient must originate at that center.

All trauma calls that meet the Trauma Field Triage Criteria Protocol, regardless of location, that are transported by air ambulance are to be transported to University Medical Center/Trauma and the medical directions for the treatment of the patient must originate at that center.

EXCEPTIONS:

1. Nothing contained within these guidelines precludes transport to any trauma facility if, in the provider's judgment, time to transport to the designated center would be unduly prolonged due to traffic and/or weather conditions and might jeopardize the patient's condition.
2. Additionally, nothing contained within these guidelines precludes transport to the closest facility if, in the provider's judgment, an ability to adequately ventilate the patient might result in increased patient mortality.

Waiting Room Criteria

Upon arrival in the emergency department, if transfer of care has not occurred in accordance with NRS 450B.790, any patient, excluding patients placed on a legal psychiatric hold, meeting *ALL* the following criteria may be placed in the hospital waiting room or other appropriate location:

1. Normal vital signs
 - A. Heart rate 60 - 100
 - B. Respiratory rate 10 - 20
 - C. Systolic BP 100 - 180
 - D. Diastolic BP 60 - 110
 - E. Room air pulse oximetry >94%
 - F. Alert and oriented x 4
2. Did not receive any parenteral medications during EMS transport except a single dose of analgesia and/or an antiemetic.
3. In the judgment of the Paramedic, does not require continuous cardiac monitoring. Note: Any ECG monitoring initiated by a transferring facility may not be discontinued by EMS personnel.
4. Can maintain a sitting position without adverse impact on their medical condition.
5. Is left with a verbal report to hospital personnel.

PROCEDURES PROTOCOLS

Cervical Stabilization

LEVEL: EMT/AEMT/Paramedic



Cervical stabilization is indicated in any patient who meets the indications (A-E) below:

Indications:

This procedure may be performed on any patient with potential for spinal injury based upon the following (NEXUS) criteria:

- A. Midline cervical spinal tenderness
- B. Focal neurologic deficit
- C. Altered mental status
- D. Evidence of drug and/or alcohol intoxication
- E. Any painful, distracting injury

Contraindications:

Cervical stabilization is *NOT* performed in the following conditions:

- A. Penetrating trauma to the head and/or neck and no evidence of spinal injury
- B. Injuries where placement of the collar might compromise patient assessment, airway management, ventilation and/or hemorrhage control
- C. Patients in cardiac arrest

Key procedural considerations:

- A. If (A-E) above are *ALL NEGATIVE*, cervical stabilization is not required.
- B. If required, cervical stabilization is the placement of an approved, properly-sized cervical collar before the patient is moved.
- C. Tape, head straps, wedges, and head and/or neck support devices are not recommended.
- D. Patients found in motor vehicles should be asked if they are able to exit the motor vehicle on their own. If so, they should be assisted to a soft stretcher and secured for transport. Patients unable to exit the vehicle on their own accord should be removed by the appropriate extrication method.
- E. Once on the stretcher, the patient may be moved to a semi-Fowler's or high-Fowler's position for comfort.
- F. If a backboard is used for extrication or movement, the patient should be immediately moved to a soft mattress, if possible.
- G. In special situations, alternate stabilization devices (e.g. vacuum mattress, KED, etc. may be used as indicated).
- H. Pediatric patients may be stabilized in an approved car seat or with a commercial pediatric stabilization device.

Continuous Positive Airway Pressure (CPAP)

LEVEL: Paramedic/AEMT

Indications:

This procedure may be performed on any patient 18 years old or older in CHF, Respiratory Distress with Bronchospasm, and pneumonia, who has *TWO* of the following:

- A. Retractions or accessory muscle use
- B. Respiratory rate >25 per minute
- C. SpO₂ ≤94%

Contraindications:

- A. Apnea
- B. Vomiting or active GI bleed
- C. Major trauma/pneumothorax
- D. Altered Mental Status

Use device per
manufacturer
instructions

Key procedural considerations:

- A. Assess patient and document VS, SpO₂ and ETCO₂ if available prior to applying oxygen. Paramedics must document ETCO₂.
- B. Select the appropriate size face mask for the patient.
- C. Inform patient about procedure process.
- D. Gradually increase the flow rate, slowly reaching the desired CPAP pressure.
- E. Secure face mask onto patient face using the head harness.
- F. Check the mask and tubing for leaks.
- G. Reassess patient and document every five minutes.
- H. If patient develops any of the contraindications or requires definitive airway control, discontinue CPAP and provide necessary airway control.

Electrical Therapy/Defibrillation

LEVEL: Paramedic

Indications:

This procedure may be performed on any patient experiencing:

- A. Ventricular fibrillation
- B. Pulseless ventricular tachycardia
- C. Torsades de Pointes

Contraindications: None

Use device per
manufacturer
instructions

Key procedural considerations:

- A. The initial and subsequent attempts shall be at the energy level(s) suggested by the device manufacturer and/or the agency's medical director.
- B. Defibrillation should be immediately provided in an arrest *WITNESSED* by EMS personnel. In an arrest that is *UNWITNESSED* by EMS personnel, two minutes of CPR should be provided prior to defibrillation.
- C. Patients with automatic implantable cardioverter-defibrillators (AICD) will need external defibrillation if the AICD is ineffective.
- D. If defibrillation is needed on a patient with a permanent implanted pacemaker, the defibrillator paddles or self adhesive electrodes should be placed at least one inch from the pulse generator of the pacemaker.

Initial attempt at pediatric defibrillation shall be at 2 J/kg. If unsuccessful, defibrillation should be attempted at 4 J/kg. Repeated defibrillations should be at >4 J/kg to 10 J/kg until conversion occurs. Adult paddles/pads may be used in children weighing more than 10 kg.

Electrical Therapy/Synchronized Cardioversion

LEVEL: Paramedic



The patient *MUST* be on a cardiac monitor and *SHOULD* have Vascular Access

Indications:

This procedure may be performed on any patient experiencing:

- A. Ventricular tachycardia with inadequate perfusion
- B. Supraventricular tachycardia with inadequate perfusion
- C. Ventricular tachycardia with adequate perfusion, but refractory to drug therapy

Adjunctive therapy:

In the conscious patient with a systolic blood pressure of >90mmHg consider:

Sedation: Etomidate 0.15 mg/kg IV/IO or;

Midazolam 0.1 mg/kg IN/IM/IV/IO, max dose 5mg. May repeat X 1 after 5 min at 0.05 mg/kg IN/IM/IV/IO, max dose 2.5 mg
Further doses with Physician Order

Or;

Diazepam 5 mg IV/IO. May repeat after five minutes with physician order.

Analgesia: Morphine Sulfate up to 0.1 mg/kg slow IV/IO to a maximum single dose of 10 mg. May repeat every 10 minutes until pain is relieved or respiratory/mental status depression occurs or;

Fentanyl up to 1 mcg/kg IN/IM/IV/IO to a maximum single dose of 100 mcg. May repeat dose after 10 minutes with physician order or;

Hydromorphone up to 1 mg IV/IO. May repeat dose after ten minutes with physician order or;

Ketamine 0.2 mg/kg IM/IV/IN/IO no repeat dose.

Contraindications: None

Key procedural considerations:

- A. Biphasic device: The initial and subsequent attempts shall be at the energy level(s) suggested by the device manufacturer and/or the agency's medical director.
- B. Monophasic device:
 - 1. Ventricular dysrhythmias: 100 J escalating to 200, 300, and 360
 - 2. Supraventricular dysrhythmias: 50 J with subsequent attempts at 100 J



Initial attempt at pediatric cardioversion shall be at 0.5 J/kg.

If unsuccessful, cardioversion should be attempted at 2 J/kg.

Adult paddle/pads may be used in children weighing more than 10 kg.

Electrical Therapy/Transcutaneous Pacing

LEVEL: Paramedic

Indications:

This procedure may be performed on any patient experiencing:

- A. Hemodynamically unstable bradycardia
- B. Unstable clinical condition that is likely because of bradycardia
- C. For pacing readiness (i.e. standby mode) in the setting of MI with bradycardia, second degree type II AV block, third degree AV block, new left or right alternating BBB or bifascicular block
- D. Overdrive pacing of tachycardias refractory to drug therapy or electrical cardioversion

Contraindications: None

Adjunctive therapy:

In the conscious patient with a systolic blood pressure of >90mmHg consider:

Sedation: Etomidate 0.15 mg/kg IV/IO or;

Midazolam 0.1 mg/kg IN/IM/IV/IO, max dose 5 mg. May repeat X 1 after 5 min at 0.05 mg/kg, max dose 2.5 mg. Further doses with Physician Order or;

Diazepam 5 mg IV/IO. May repeat after five minutes with physician order.

Analgesia: Morphine Sulfate up to 0.1 mg/kg slow IV/IO to a maximum single dose of 10 mg. May repeat every 10 minutes until pain is relieved or respiratory/mental status depression occurs or;

Fentanyl up to 1 mcg/kg IN/IM/IV/IO to a maximum single dose of 100 mcg. May repeat dose after 10 minutes with physician order or;

Hydromorphone up to 1 mg IV/IO. May repeat dose after ten minutes with physician order or;

Ketamine 0.2 mg/kg IM/IV/IN/IO no repeat dose.

Key procedural considerations:

- A. Apply pacing pads, begin pacing at a rate of 60 beats per minute at the lowest available current.
- B. Increase current by 20 milliamp increments until electrical capture.
- C. In the event of electrical capture and no pulses, continue pacing and CPR.



Pediatric pacing is by telemetry physician order only

Endotracheal Intubation

LEVEL: Paramedic

1. All intubations **MUST** have initial, en route, and at transfer of care End-Tidal CO₂ detection/capnography performed and recorded on the PCR.
2. All intubation attempts **MUST** be documented on the PCR.

Indications:

This procedure may be performed on any patient in whom attempts at basic airway and ventilatory support are unsuccessful **AND** who has at least one of the following:

- A. Hypoxia
- B. Respiratory arrest/failure
- C. Inability to maintain airway patency

Contraindications:

Absolute Contraindications: None

Relative Contraindications:

- A. Presence of gag reflex
- B. Suspected narcotic overdose/hypoglycemia prior to administration of Naloxone/Glucose

Induction:

Etomidate 0.3 mg/kg IV/IO, max dose 30 mg. Pediatric patients have a max dose of 20 mg.

OR

Ketamine 2 mg/kg IV/IO or 4 mg/kg IM.

OR



Midazolam (pediatric patients only) 0.1 mg/kg IV/IO/IN, max single dose 5 mg. Needs to be given slowly over 5 min. Repeat doses with physician order only.

Post-Intubation Sedation:



***Note:** Sedation should be titrated for patient comfort and respiratory compliance. Always use the least amount of medication possible to maintain sedation effects and reduce patient agitation*

Midazolam 0.1 mg/kg IV/IN, max dose 5 mg. May repeat X1 after 5 min at 0.05 mg/kg IV/IN/IM, max dose 2.5 mg. Repeat doses with physician order only.

OR

Ketamine 2 mg/kg IV or 4 mg/kg IM. Repeat doses with physician order only.

Post-Intubation Analgesia:



Fentanyl 1 mcg/kg IV/IM/IN/IO, max dose 100 mcg.

Endotracheal Intubation (Cont.)

LEVEL: Paramedic

Key procedural considerations:

- A. Position head properly.
- B. Insert blade while displacing tongue and elevate mandible with laryngoscope.
- C. Introduce ET tube and advance to proper depth.
- D. Inflate cuff to proper pressure and disconnect syringe.
- E. Ventilate patient and confirm proper placement.
- F. Verify proper tube placement by secondary confirmation such as capnography or colorimetric device.
- G. Secure device or confirm that the device remains properly secured.

Nasotracheal Intubation:

Contraindications:

- A. Apnea or near-apnea
- B. Suspected basilar skull, nasal, or midface fractures
- C. Coumadin anticoagulation therapy or hemostatic disorders
- D. Upper neck hematomas
- E. Should *NOT* be attempted in children

Adjunctive Therapy:

Prep the nostrils with:

- 1) Phenylephrine 2-3 drops (or 1-2 sprays in each nostril),
or
Oxymetazoline 2 sprays in each nostril.
and
- 2) Lidocaine 2% lubricant.

Check and prepare the
endotracheal airway device prior
to insertion

Key procedural considerations:

- A. Position patient semi-Fowler, sitting or supine.
- B. Insert lubricated ET tube into dilated nostril and advance straight back (posteriorly).
- C. Listen to end ET tube for sounds of patient's breathing.
- D. During inhalation, smoothly advance tube through glottic opening.
- E. Inflate cuff to proper pressure and disconnect syringe.
- F. Verify proper tube placement by secondary confirmation such as capnography or colorimetric device.
- G. Secure device or confirm that the device remains properly secured.

Extraglottic Airway Device

LEVEL: AEMT/Paramedic

Indications:

This procedure may be performed on any patient in which attempts at basic airway and ventilatory support are unsuccessful *AND* who has at least one of the following:

- A. Hypoxia
- B. Respiratory arrest/failure
- C. Obtundation
- D. Failed endotracheal intubation

Contraindications:

- A. Gag reflex
- B. History of esophageal trauma, or known esophageal disease
- C. Recent ingestion of a caustic substance
- D. Tracheostomy or laryngectomy
- E. Suspected foreign body obstruction

Check and prepare the
extraglottic airway
device prior to insertion

Key procedural considerations:

- A. Pre-oxygenate the patient.
- B. Position the patient's head in a neutral or slightly flexed position if no suspected spinal injury (if a spine injury is suspected, maintain a neutral, in-line head position).
- C. Perform a tongue-jaw lift.
- D. Insert device to proper depth. *NEVER* force. If device does not advance, readjust the insertion.
- E. Secure device in the patient (inflate cuff(s) with proper volume(s) and immediately remove syringe).
- F. Ventilate patient and confirm proper ventilation (correct lumen and proper insertion depth) by auscultation bilaterally over lungs and over epigastrium.
- G. Adjust ventilation as necessary (ventilate through additional lumen or slightly withdraw tube until ventilation is optimized).
- H. Verify proper tube placement by secondary confirmation such as capnography or colorimetric device.
- I. Secure device or confirm that the device remains properly secured.

First Response Evaluate/Release

LEVEL: AEMT/PARAMEDIC

Inclusion Criteria:

- A. Coded and dispatched using MPDS as an Alpha or Omega category
- B. Patient age $\geq 18 \leq 65$
- C. Full assessment performed by first response
- D. Patient deemed to have decision making capacity
- E. Normal vital signs including SPO₂
- F. Patient has a phone, ability and willingness to call 9-1-1 if their condition worsens
- G. In the opinion of the AEMT/paramedic and the patient it is safe to release until an ambulance arrives

Exclusion Criteria:

- A. Abnormal vital signs including SPO₂
- B. Pregnancy
- C. Any high risk complaints/symptoms
 - a. Chest pain
 - b. Signs/symptoms of possible stroke
 - c. Allergic reaction
 - d. Shortness of breath
 - e. Abdominal pain/flank pain above umbilicus age >35
 - f. Syncope, near syncope, dizziness
 - g. Seizure
 - h. History or sign of head trauma
 - i. Active bleeding
 - j. Threat to self or others
 - k. Overdose or ingestional error
 - l. Patients meets Trauma Field Triage Criteria
- D. No SNHD EMS Protocol indication for obtaining EKG or placing the patient on a cardiac monitor

EMS patient care record must be completed within four hours of clearing the call

The Field Response Low-Risk Alpha Evaluate and Release Form must be completed and a copy left with the patient for inclusion in the secondary responder's patient care report

Hemorrhage Control

LEVEL: EMT/AEMT/Paramedic

Hemorrhage:

This procedure may be performed on any patient that has bleeding from an extremity, junctional hemorrhage or torso hemorrhage.


Extremity Hemorrhage – Tourniquet Application:

- A. Apply tourniquet proximal to the bleeding site.
- B. Absolute contraindication: Bleeding has stopped
- C. If bleeding is not controlled, consider additional tightening or applying a second tourniquet proximal side by side to the first.
- D. Wound packing does not preclude you from placing a tourniquet.


Junctional Hemorrhage – Wound Packing:

- A. Junctional Hemorrhage Defined: hemorrhage occurring at the junction of an extremity with the torso, and/or the base of the neck.
- B. Use direct pressure and an appropriate pressure dressing with deep wound packing (plain gauze or, if available, hemostatic gauze).
- C. Absolute Contraindication: Hemostatic gauze use on hemorrhaging abdominal wounds.

Tourniquet Placement



Junctional Hemorrhage



Medication Administration

LEVEL: EMT/AEMT/Paramedic (based on medication)

Indications:

This procedure may be performed on any patient that requires the administration of a medication.

Key procedural considerations (GENERAL):

- A. Inquire about allergies and previous medication reactions
- B. Check and recheck medication
- C. Solution clarity and expiration date
- D. Right drug
 - Right patient
 - Right dose
 - Right time
 - Right route
 - Right documentation
- E. Dispose of syringe and other material in proper container

Intravenous and Intraosseous Bolus Medications

Key procedural considerations:

- A. Identify and cleanse injection site closest to the patient
- B. Administer correct dose at proper push rate
- C. Turn IV on and adjust drip rate to TKO/KVO

Intramuscular and Subcutaneous Drug Administration

Key procedural considerations:

- A. Needle should be 20 gauge or smaller
- B. Locate administration site
 - Deltoid muscle
 - Vastus lateralis (lateral thigh) muscle
 - Ventrogluteal or dorsogluteal muscles (buttocks)

IM

Pull skin tight
Insert needle at a 90° angle to the skin
Advance into muscle layer

SQ

Pinch to lift skin slightly
Insert needle at a 45° angle to the skin
Advance into subcutaneous layer

Medication Administration (Cont.)

LEVEL: EMT/AEMT/Paramedic (based on medication)

Mucosal Atomizer Device (MAD) Administration

Medications: Fentanyl, Ketamine, Midazolam, Naloxone Hydrochloride

Key procedural considerations:

- A. Using the free hand, hold the crown of the head stable.
- B. Place the tip of the MAD snugly against the nostril, aiming slightly up and outward (toward the top of the ear).
- C. Briskly compress the syringe to deliver half the medication into the nostril.
- D. Move the device over to the opposite nostril and administer the remaining medication.

Intravenous and Intraosseous Bolus Medications

Key procedural considerations:

- A. Identify and cleanse injection site closest to the patient
- B. Administer correct dose at proper push rate
- C. Turn IV on and adjust drip rate to TKO/KVO

Indications:

This procedure may be performed on any patient that requires the administration of a medication.

Key procedural considerations (GENERAL):

- A. Inquire about allergies and previous medication reactions
- B. Check and recheck medication
- C. Solution clarity and expiration date
- D. Right drug
 - Right patient
 - Right dose
 - Right time
 - Right route
 - Right documentation
- E. Dispose of syringe and other material in proper container

Needle Cricothyroidotomy

LEVEL: Paramedic

Indications:

This procedure may be performed on any patient with:

- A. Total airway obstruction by any BLS or ALS procedures, *OR*
- B. Inability to be adequately ventilate with any provider level emergency care procedures prior to the attempt.

Contraindications:

- A. Inability to identify landmarks (cricothyroid membrane)
- B. Underlying anatomical abnormality (tumor)
- C. Tracheal transection
- D. Acute laryngeal disease due to infection or trauma



- Pediatric needle cricothyroidotomy is by Telemetry Physician order only.
- You **MUST** use a 14 gauge over-the-needle catheter attached to a 10 cc syringe or commercial cricothyroidotomy device.

Key procedural considerations:

Please follow Manufacturer's Instructions

- A. Position patient supine (if possible), hyperextending the head.
- B. Locate cricothyroid membrane and clean site thoroughly.
- C. Stabilize cricoid and thyroid cartilages with one hand.
- D. Puncture needle/catheter at a 90° angle and then change insertion angle to 45° up to the stopper; gently aspirate with attached syringe.
- E. When syringe is able to aspirate air, stop advancing needle.
- F. Remove the stopper from the cannula and advance the cannula only until the phlange is flush with the patient's neck. Remove the metal needle from the cannula. Remove the syringe.
- G. Secure the cannula with the neck strap.
- H. Apply connecting tube and attach to BVM and ventilate patient.

Needle Thoracentesis

LEVEL: Paramedic

Indications:

This procedure may be performed on any patient who has evidence of a tension pneumothorax, demonstrated by the presence of:

Progressive respiratory distress and/or increased resistance to bagging, *AND* unilateral diminished/absent breath sounds, associated with:

- A. Tracheal deviation, or
- B. Jugular venous distension, or
- C. Signs of shock, low BP with chest trauma present

Contraindications: None



Needle Thoracentesis is permitted in pediatric patients.

Key procedural considerations:

- A. Primary site is the 2nd intercostal space mid-clavicular line of the affected side.
- B. Alternate location is the 4th - 5th intercostal space in the mid-axillary line of the affected side.
- C. Use a site specific, appropriate length needle to decompress the chest.
- D. Prep site with iodine and/or alcohol.
- E. Place tip of needle on top of appropriate rib and insert over top of rib into intercostal space.
- F. Advance catheter and remove needle.
- G. Secure catheter and consider attaching a flutter valve assembly.

Tracheostomy Tube Replacement

LEVEL: Paramedic

Indications:

This procedure may be performed on any patient that has A *TRACHEOSTOMY TUBE* and *WHO HAS*:

- A. Hypoxia
- B. Respiratory arrest/failure
- C. Obtundation
- D. Secretions unable to be cleared by suctioning

Contraindications: None

Key procedural considerations:

- A. If the patient or family has a replacement tube available, it may be used. If a replacement tube is not available, an endotracheal tube of a similar outer diameter may be used.
- B. Premoisten the tube with water soluble lubricant.
- C. Extend the neck and, if necessary, place a roll between the patient's shoulder blades to aid in visualizing the stoma.
- D. If the tube cannot be placed easily, withdraw the tube; administer oxygen and positive pressure ventilation. *NEVER* force the tube.
- E. Secure the device to the patient.
- F. If the tube cannot be easily placed, a suction catheter may be used as a guide.

Traction Splint

LEVEL: EMT/AEMT/Paramedic

Indications:

This procedure may be performed on any patient with an isolated midshaft femur fracture.

Contraindications:

- A. Pelvic fracture or instability**
- B. Knee, lower leg, or ankle instability**

Key procedural considerations:

- A. Assess motor, sensory, and circulatory function in the involved extremity.**
- B. Apply traction splint per the manufacturer's guidelines.**
- C. Initiate mechanical traction to match manual traction.**
- D. Reassess motor, sensory, and circulatory function in the involved extremity.**
- E. Exercise care when applying traction not to reintroduce bone ends into the body.**

Vagal Maneuvers

LEVEL: Paramedic



The patient ***MUST*** be attached to a cardiac monitor and ***MUST*** have vascular access prior to performing the procedure

Indications:

This procedure may be performed on any patient who is experiencing Supraventricular Tachycardia with adequate perfusion.

Contraindications:

None

Key procedural considerations:

A. Approved methods include:

1. Valsalva maneuver
2. Head-down tilt with deep inspiration
3. Activation of the “diving reflex” by facial immersion in ice water (unless ischemic heart disease is present)
4. Carotid massage (only on patients under 40 years of age)



B. In infants and young children, the most effective vagal maneuver is the application of ice to the face. IV access is not mandatory prior to vagal maneuvers in children.

Vascular Access

LEVEL: AEMT/Paramedic



Vascular access attempts should not unnecessarily delay transport:
Attempts should be completed en route.
All attempts are to be documented on the PCR.

Indications for Peripheral Vascular Access:

This procedure may be performed on any patient whenever there is a potential need for:

- A. Intravenous drug administration
- B. Need to administer IV fluids for volume expansion

Contraindications: None

Key procedural considerations:

- A. Saline locks may be used when appropriate and flushed with a 3 cc bolus of NS as needed.
- B. Extension tubing should be used on all IV lines.

Indications for Intraosseous Access (Paramedic for Adult and Peds, AEMT for Adult Only)

Critically ill or injured patient who requires IV drugs/fluids and in whom a peripheral line cannot be immediately established.

Contraindications:

- A. Placement in, or distal to a fractured bone.
- B. Previous significant orthopedic procedure at the site; prosthetic limb or joint; IO catheter use in past 48 hours of the target bone.
- C. Infection at the area of insertion.
- D. Absence of adequate anatomical landmarks.

Paramedic may administer lidocaine 1% or 2% preservative-free for anesthetic in a patient responsive to pain.

- 1) Prime IO extension tubing set with lidocaine (EZ IO, EZ Connect priming volume is 1ml)
- 2) Slowly infuse lidocaine 40mg (PEDIATRIC dose: 0.5 mg/kg not to exceed 40 mg) IO over 120 seconds.
- 3) Allow lidocaine to dwell in IO space for 60 seconds.
- 4) Flush IO with 5-10 ml normal saline.
- 5) Slowly administer an additional dose of lidocaine IO (20mg) over 60 seconds. (Pediatric dose: 0.25 mg/kg, not to exceed 20 mg)
- 6) Consider systemic pain medication for patients not responding to IO lidocaine.

Key procedural considerations: Only 1 (one) attempt is permitted per extremity

Indications for use of Previously Established Central Line Access:

This procedure may be performed on any critically ill or injured patient who requires IV drugs or IV fluids AND in whom a peripheral line cannot be established.

Contraindications: Inability to freely aspirate blood out of the catheter.

Key procedural considerations: Central line access (Implantable Ports, Port-A-Caths, Medports)



- A. May only be used if the device has already been accessed and IV fluid set-up has been established and running.
- B. These devices require special needles (non-coring type) for access. The device may be damaged if standard jumper (conventional) needles are used to access the ports.

FORMULARY

ACETYLSALICYLIC ACID (Aspirin)

CLASS: Nonsteroidal anti-inflammatory (NSAID)

ACTION: Platelet inhibition

DOSE:

Protocol	PO
Chest Pain (Non Traumatic) and Suspected ACS (29)	324 mg (81 mg tablet X 4) chew and swallow
STEMI (Suspected) (61)	324 mg (81 mg tablet X 4) chew and swallow

CONTRAINDICATIONS: Allergy to Aspirin

ADVERSE REACTIONS: None

ADENOSINE (Adenocard)

CLASS: Antiarrhythmic

ACTION: Slows conduction through the AV node and can interrupt re-entry pathways

DOSE:

Protocol	IV/IO	Repeat?	Notes
Tachycardia/Stable (65)	6 mg Rapid IVP/IO	If unsuccessful, repeat with 12 mg rapid IVP/IO	
Tachycardia/Unstable (67)	12 mg rapid IVP/IO	No	
Pediatric Tachycardia/Stable (112)	0.1mg/kg rapid IVP/IO, not to exceed 6 mg	0.2mg/kg rapid IVP/IO, not to exceed 12 mg	Initial and repeat doses by telemetry order only
Pediatric Tachycardia/Unstable (114)	0.2mg/kg rapid IVP/IO, not to exceed 12 mg	No	Initial dose by telemetry order only

CONTRAINDICATIONS: Second or third degree AV block or sick sinus syndrome (unless patient with a functional artificial pacemaker); atrial flutter; atrial fibrillation

Repeat doses of Adenosine are not indicated if the dysrhythmia reoccurs after conversion. Alternate pharmacological therapy may be necessary.

ADVERSE REACTIONS: facial flushing; headache; sweating; palpitations

ALBUTEROL (Proventil)

CLASS: Sympathomimetic

ACTION: Bronchodilator

DOSE:

Protocol	Nebulized	Repeat
Allergic Reaction (17)	2.5 mg in 3 ml SVN	Yes until improved
Drowning (35)	2.5 mg in 3 ml SVN	Yes until improved
Hyperkalemia (41)	2.5 mg in 3 ml SVN	Continuous
Respiratory Distress (51)	2.5 mg in 3 ml SVN	Yes until improved
Pediatric Allergic Reaction (78)	2.5 mg in 3 ml SVN	Yes until improved
Pediatric Drowning (92)	2.5 mg in 3 ml SVN	Yes until improved
Pediatric Respiratory Distress (104)	2.5 mg in 3 ml SVN	Yes until improved

CONTRAINDICATIONS: Hypersensitivity to the drug

ADVERSE REACTIONS: Tachycardia; palpitations; anxiousness; headache

AMIODARONE (Cordarone)

CLASS: Antiarrhythmic

ACTION: Suppresses ventricular ectopy; increases ventricular fibrillation threshold

DOSE:

Protocol	IV/IO	Repeat	Note
Cardiac Arrest (27)	300 mg IV/IO	150 mg if refractory after 5th shock	Consider H's & T's
Tachycardia/Stable (65)	150 mg in 50 cc NS over 10 min	No	By Physician Order Only
Tachycardia/Unstable (67)	150 mg in 50 cc NS over 10 min	No	
Pediatric Cardiac Arrest (88)	5 mg/kg IV/IO	5 mg/kg IV/IO if refractory after 5th shock	Consider H's & T's
Peds Tachycardia/Stable (112)	5 mg/kg in 50 cc NS over 20 min	No	
Peds Tachycardia/Unstable (114)	5 mg/kg IV/IO in 50 cc NS over 20 min	No	

CONTRAINDICATIONS: Hypersensitivity to the drug; cardiogenic shock; high grade AV block; marked sinus bradycardia; or bradycardia with ventricular escape beats

ADVERSE REACTIONS: Seizures; respiratory depression; dizziness; restlessness; confusion; tinnitus; blurred vision; numbness; muscle twitching; hypotension; bradycardia; heart block; nausea; vomiting

ATROPINE SULFATE

CLASS: Parasympathetic blocker

ACTION: Cholinergic blocking agent; increases rate of SA node discharge; increases conduction

DOSE:

Protocol	IV/IO	Repeat	Note
Bradycardia (23)	0.5 mg IV/IO	Yes, q 5 min to max 3 mg	
Overdose/Poisoning (45)	2 mg IV/IO	Yes, q 15 min as needed	For organophosphate toxicity. Administer to decrease secretions and ventilatory resistance
Pediatric Bradycardia (84)	0.02mg/kg IV/IO, min dose 0.1mg, max dose 0.5mg	Yes, X1 after 5 min	
Peds Overdose/Poisoning (100)	0.02mg/kg IV/IO, min dose 0.1mg	Yes, q 15 min as needed	For organophosphate toxicity. Administer to decrease secretions and ventilatory resistance

CONTRAINDICATIONS: none

ADVERSE REACTIONS: none

BRONCHODILATOR METERED DOSE INHALER

CLASS: Sympathomimetic

ACTION: Bronchodilator

DOSE: Assist the patient in administering his or her own Bronchodilator Metered Dose Inhaler exactly as prescribed.

CONTRAINDICATIONS: Sensitivity to the drug

ADVERSE REACTIONS: Tachycardia; palpitations; anxiousness; headache

CALCIUM CHLORIDE

CLASS: Electrolyte

ACTION: Increases myocardial contractility; increases myocardial excitability; decreases heart rate

CALCIUM CHLORIDE (continued)

DOSE:

Protocol	IV/IO	Note
Bradycardia (23)	1 g IV/IO	For patients on Calcium Channel blockers
Hyperkalemia (41)	1 g slow IVP/IO	Contraindicated in patients taking digatalis products
Overdose/Poisoning (45)	1 g slow IVP/IO	For calcium channel blocker overdose
Peds Overdose/Poisoning (100)	20 mg/kg slow IVP/IO	For calcium channel blocker overdose

CONTRAINDICATIONS: Patients receiving digitalis

ADVERSE REACTIONS: None

DIAZEPAM (Valium)

CLASS: Antianxiety/Anticonvulsant

ACTION: CNS Depressant

DOSE:

Protocol	IV/IO/IM	PR	Repeat?
Behavioral Emergency (21)	5 mg IV/IO		Yes, q 5 min
Obstetrical Emergency (43)	5 mg IV		Yes, X1. Orders needed for additional
Seizure (53)	5 mg IV/IM/IO		
Ventilation Management (69)	5 mg IV/IO		With Physician order only
Pediatric Seizure (106)	0.1mg/kg IV/IM/IO, max dose 5 mg	0.1 mg/kg, max dose 5 mg	No
Peds Ventilation Management (116)	0.2mg/kg IV/IO, max dose 5 mg	0.5 mg PR, max dose 20 mg	With Physician order only
Electrical Therapy/Synchronized Cardioversion (140)	5 mg IV/IO		With physician order only
Electrical Therapy/ Transcutaneous Pacing (141)	5.mg IV/IO		With Physician order only

CONTRAINDICATIONS: Hypersensitivity; hypotension

ADVERSE REACTIONS: Respiratory depression; CNS depression; nausea; vomiting

DIPHENHYDRAMINE HYDROCHLORIDE (Benadryl)

CLASS: Antihistamine

ACTION: Blocks histamine receptors; has some sedative effects; anticholinergic

DOSE:

Protocol	IM/IV/IO	PO
Allergic Reaction (17)	50 mg IM/IV/IO	50mg PO
Overdose/Poisoning (45)	50 mg IV/IM/IO	
Peds Allergic Reaction (78)	1 mg/kg IV/IM/IO, max 50 mg	less than 6 years: 12.5 mg PO; 6 to 12 years: 25 mg PO; >12 years: 50 mg PO
Peds Overdose/Poisoning (100)	1 mg/kg IV/IM/IO, max 50 mg	

CONTRAINDICATIONS: Hypersensitivity to the drug

ADVERSE REACTIONS: Sedation; palpitations; decreased blood pressure; headache; dries (thickens) bronchial secretions; blurred vision

DOPAMINE HYDROCHLORIDE (Intropin)

CLASS: Sympathomimetic

ACTION: Positive inotrope with dose-related vascular effects

DOSE:

Protocol	IV/IO
Allergic Reaction (17)	5-20 mcg/kg/min IV/IO, titrate to SBP >90
Bradycardia (23)	5-10 mcg/kg/min IV/IO, titrate to SBP >90, max dose 20mcg/kg/min
Pulmonary Edema/CHF (49)	5-20 mcg/kg/min IV/IO, titrate to SBP >90
Sepsis (55)	5-20 mcg/kg/min IV/IO, titrate to SBP >90
Shock (57)	5-20 mcg/kg/min IV/IO, titrate to SBP >90
Pediatric Shock (108)	5-20 mcg/kg/min IV/IO, titrate to SBP >70 + 2X age

DOPAMINE HYDROCHLORIDE (continued)

CONTRAINDICATIONS: Hypovolemic shock is a relative contraindication. Hypotension due to hypovolemia or distributive shock should be addressed with a fluid bolus before administering Dopamine.

ADVERSE REACTIONS: Ventricular tachycardia; ectopic beats; nausea; vomiting; dyspnea; hypertension and extreme vasoconstriction may occur with high infusion rates; hypotension may occur with low infusion rates

DROPERIDOL (Inapsine)

CLASS: Antiemetic

ACTION: Lowers incidence of nausea and vomiting

DOSE:

Protocol	IM/IV/IO	Repeat?
Abdominal/Flank Pain, Nausea & Vomiting (15)	1.25 mg IM/IV/IO	
Behavioral Emergency (21)	1.25 mg IM/IV/IO, followed by saline flush or bolus	Yes, q 5 min
Pain Management (47)	1.25 mg IM/IV/IO	

CONTRAINDICATIONS: Contraindicated for any patient having suspected STEMI and chest pain; hypotensive patients; respiratory depression; known hypersensitivity to the drug; known prolonged QT interval

ADVERSE REACTIONS: EPS; syncope; cardiac dysrhythmias

EPINEPHRINE 1:1000

CLASS: Sympathomimetic

ACTION: Bronchodilation; positive chronotrope; positive inotrope

DOSE:

Protocol	IM/ETT/SVN	Repeat?
Allergic Reaction (17)	0.5 mg IM	yes, q 15 min to a max of 1.5 mg
Peds Allergic Reaction (78)	0.01 mg/kg IM, max single dose 0.5 mg	Yes, q 5 min to a max of 1.5 mg
Peds Bradycardia (84)	0.1 mg/kg ETT max 1 mg	Yes, q 3-5 min
Peds Cardiac Arrest (88)	0.1 mg/kg ETT max 1 mg	Yes, q 3-5 min
Peds Respiratory Distress (104)	3-5 mg SVN if SPO2 <94%	

CONTRAINDICATIONS: None

EPINEPHRINE 1:1000 (continued)

ADVERSE REACTIONS: Palpitations due to tachycardia or ectopic beats may produce arrhythmia if cardiac disease is present; elevation of blood pressure; headache; anxiousness

EPINEPHRINE 1:10,000

CLASS: Sympathomimetic

ACTION: Bronchodilation; positive chronotrope; positive inotrope

DOSE:

Protocol	IV/IO	ETT	Repeat?
Cardiac Arrest (27)	1 mg IV/IO	2-2.5 mg ETT	Yes, q 3-5 min
Peds Bradycardia (84)	0.01 mg/kg IV/IO		Yes, q 3-5 min
Peds Cardiac Arrest (88)	0.01 mg/kg IV/IO		Yes, q 3-5 min
Neonatal Resuscitation (98)	0.01 mg/kg IV/IO		As needed every 3-5 min for HR <60

CONTRAINDICATIONS: None

ADVERSE REACTIONS: Palpitations due to tachycardia or ectopic beats may produce arrhythmia if cardiac disease is present; elevation of blood pressure; headache; anxiousness

EPINEPHRINE 1:100,000

CLASS: Sympathomimetic

ACTION: Bronchodilation; positive chronotrope; positive inotrope

DOSE:

Protocol	IV/IO	Repeat?	Preparation
Allergic Reaction (17)	Push Dose 1:100,000 5-10 mcg IV/IO	Yes, q 2-5 min to maintain SBP >90	To Prepare: mix cardiac epinephrine 1:10,000 1 ml PLUS 9 ml Normal Saline = 10 ml EPINEPHRINE 1:100,000 at 10 mcg/ml
Sepsis (55)	Push Dose 1:100,000 5-10 mcg IV/IO	Yes, q 2-5 min to maintain SBP >90	To Prepare: mix cardiac epinephrine 1:10,000 1 ml PLUS 9 ml Normal Saline = 10 ml EPINEPHRINE 1:100,000 at 10 mcg/ml
Shock (57)	Push Dose 1:100,000 5-10 mcg IV/IO	Yes, q 2-5 min to maintain SBP >90	To Prepare: mix cardiac epinephrine 1:10,000 1 ml PLUS 9 ml Normal Saline = 10 ml EPINEPHRINE 1:100,000 at 10 mcg/ml

CONTRAINDICATIONS: None

ADVERSE REACTIONS: Palpitations due to tachycardia or ectopic beats may produce arrhythmia if cardiac disease is present; elevation of blood pressure; headache; anxiousness

EPINEPHRINE AUTO-INJECTOR

CLASS: Sympathomimetic

ACTION: Bronchodilation; positive chronotrope; positive inotrope

DOSE: Assist the patient in administering his or her own Epinephrine auto-injector exactly as prescribed

CONTRAINDICATIONS: None

ADVERSE REACTIONS: Palpitations due to tachycardia or ectopic beats may produce arrhythmia if cardiac disease is present; elevation of blood pressure; headache; anxiousness

ETOMIDATE (Amidate)

CLASS: Sedative/Hypnotic

ACTION: CNS depressant

DOSE:

Protocol	IV/IO	Notes
Tachycardia/Stable (65)	0.15 mg/kg IV/IO	Need Physician Orders
Tachycardia/Unstable (67)	0.15 mg/kg IV/IO	
Ventilation Management (69)	0.3 mg/kg IV/IO	Max dose 30 mg
Peds Tachycardia/Stable (112)	0.15 mg/kg IV/IO	Need Physician Orders
Peds Tachycardia/Unstable (114)	0.15 mg/kg IV/IO	
Peds Ventilation Management (116)	0.3 mg/kg IV/IO	Max dose for peds 20 mg
Electrical Therapy/Synchronized Cardioversion (140)	0.15 mg/kg IV/IO	
Electrical Therapy/ Transcutaneous Pacing (141)	0.15 mg/kg IV/IO	
Endotracheal Intubation (142)	0.3 mg/kg IV/IO	Max dose 20 mg for peds, 30 mg for adults

CONTRAINDICATIONS: Known hypersensitivity to the drug

ADVERSE REACTIONS: Pain; transient skeletal movements; nausea; vomiting; hypoventilation; hypotension

FENTANYL CITRATE

CLASS: Analgesic

ACTION: CNS Depressant

FENTANYL CITRATE (continued)

DOSE:

Protocol	IN/IM/IV/IO	Repeat?
Pain Management (47)	1 mcg/kg IN/IM/IV/IO, max single dose 100 mcg	Yes, X1 after 10 min
Ventilation Management (69)	1 mcg/kg IN/IM/IV/IO, max single dose 100 mcg	Need Physician Orders
Peds Pain Management (102)	1 mcg/kg IN/IM/IV/IO, max single dose 100 mcg	Need Physician Orders
Peds Ventilation Management (116)	1 mcg/kg IN/IM/IV/IO, max single dose 100 mcg	Need Physician Orders
Electrical Therapy/Synchronized Cardioversion (140)	1 mcg/kg IN/IM/IV/IO, max single dose 100 mcg	Need Physician Orders
Electrical Therapy/ Transcutaneous Pacing (141)	1 mcg/kg IN/IM/IV/IO, max single dose 100 mcg	Need Physician Orders
Endotracheal Intubation (142)	1 mcg/kg IN/IM/IV/IO, max single dose 100 mcg	Need Physician Orders

CONTRAINDICATIONS: Known hypersensitivity to the drug

ADVERSE REACTIONS: Respiratory depression; rapid infusion may produce “stiff chest syndrome”

GLUCAGON

CLASS: Insulin Antagonist

ACTION: Reverses the effects of hypoglycemia

DOSE:

Protocol	IV/IM/IO	Notes
Altered Mental Status/Syncope (19)	1 mg IM	
Bradycardia (23)	1 mg IV	For patients on beta blockers
Overdose/Poisoning (45)	1 mg IV/IM/IO	For beta blocker overdose, may repeat X1
Seizure (53)	1 mg IM	
Peds Altered Mental Status (80)	0.5 mg IM if pt is <20 kg; 1 mg IM if pt is >20 kg	
Peds Overdose/Poisoning (100)	0.5 mg IV/IM/IO	For beta blocker overdose, may repeat X1
Peds Seizure (106)	0.5 mg IM	
Peds Shock (10)	0.5 mg IM if pt is <20 kg; 1 mg IM if pt is >20 kg	

CONTRAINDICATIONS: None

ADVERSE REACTIONS: Nausea; vomiting

GLUCOSE – ORAL GLUCOSE

CLASS: Carbohydrate

ACTION: Quick infusion of sugar into blood for metabolism

DOSE:

Protocol	PO	Notes
Altered Mental Status/Syncope (19)	15 g PO	Use if patient is able to protect their own airway
Peds Altered Mental Status/Syncope (80)	Up to 15 g PO	Use if patient is able to protect their own airway
Peds Shock (108)	Up to 15 g PO	Use if patient is able to protect their own airway

CONTRAINDICATIONS: None

ADVERSE REACTIONS: None

GLUCOSE – D10

CLASS: Carbohydrate

ACTION: Quick infusion of sugar into blood for metabolism

DOSE:

Protocol	IV/IO	Repeat?
Altered Mental Status/Syncope (19)	25 g IV/IO (250 ml of 10% solution)	Yes, X1 in 5 min
Seizure (53)	25 g IV/IO (250 ml of 10% solution)	Yes, X1 in 5 min
Peds Altered Mental Status/Syncope (80)	5 ml/kg IV/IO, max single dose 25 g	
Peds Seizure (106)	5 ml/kg IV/IO, max single dose 25 g	
Peds Shock (108)	5 ml/kg IV/IO, max single dose 25 g	

CONTRAINDICATIONS: None

ADVERSE REACTIONS: None

HYDROMORPHONE (Dilaudid)

CLASS: Analgesic

ACTION: CNS Depressant

DOSE:

Protocol	IM/IV/IO	Repeat?
Pain Management (47)	0.01 mg/kg IM/IV/IO, max single dose 1 mg	Yes, X1 after 10 min
Electrical Therapy/Synchronized Cardioversion (140)	Up to 1 mg IV/IO	Need physician order
Electrical Therapy/Transcutaneous Pacing (141)	Up to 1 mg IV/IO	Need physician order

CONTRAINDICATIONS: Known hypersensitivity; intolerance to opiate analgesics

ADVERSE REACTIONS: Respiratory depression

HYDROXOCOBALAMIN

CLASS: Detoxifying agent

ACTION: Competitively binds to cyanide ions

DOSE:

Protocol	IV/IO	Notes
Overdose/Poisoning (45)	5 g IV/IO over 15 min	For cyanide poisoning
Smoke Inhalation (59)	5 g IV/IO over 15 min	
Peds Overdose/Poisoning (100)		refer to protocol
Peds Smoke Inhalation (110)		refer to protocol

CONTRAINDICATIONS: None

ADVERSE REACTIONS: None

IPRATROPIUM BROMIDE (Atrovent)

CLASS: Anticholinergic

ACTION: Appears to inhibit vagally mediated reflexes

IPRATROPIUM BROMIDE (continued)

DOSE:

Protocol	SVN	Repeat?
Respiratory Distress (51)	2.5 ml 0.02% solution SVN	No
Peds Respiratory Distress (104)	2.5 ml 0.02% solution SVN	No

CONTRAINDICATIONS: Hypersensitivity to ipratropium

ADVERSE REACTIONS: Headache; nausea

IPRATROPIUM BROMIDE and ALBUTEROL SULFATE (Duoneb)

CLASS: Anticholinergic/Sympathomimetic

ACTION: Appears to inhibit vagally mediated reflexes and acts as a bronchodilator

DOSE:

Protocol	SVN	Repeat?
Respiratory Distress (51)	3 ml SVN	No
Peds Respiratory Distress (104)	3 ml SVN	No

CONTRAINDICATIONS: Hypersensitivity to either of the base medications

ADVERSE REACTIONS: Tachycardia; palpitations; anxiousness; headache; nausea

KETAMINE (Ketalar)

CLASS: General anesthetic/ induction agent

ACTION: CNS Depressant

DOSE:

Low-Dose Protocol	IV/IO/IM/IN		Repeat OK?	Note
Pain Management (47)	0.2 mg/kg		No	Avoid in Chest Pain/ACS/STEMI
Electrical Cardioversion (140)	0.2 mg/kg		No	Given in addition to sedation
Transcutaneous Pacing(141)	0.2 mg/kg		No	Given in addition to sedation
Protocol	IV/IO	IM	Repeat OK?	
Behavioral Emergency (21)	2 mg/kg	2-4 mg/kg	No	
Ventilation Mgmt (69/116)	2 mg/kg	4 mg/kg	No	
Endotracheal Intubation (142)	2 mg/kg	4 mg/kg	Yes (X1)	

CONTRAINDICATIONS: Use caution in patients with systolic over 180 mm Hg

ADVERSE REACTIONS: Respiratory depression

LIDOCAINE (Xylocaine) 1% or 2% INJECTION

CLASS: Anesthetic

ACTION: Produces anesthesia by interfering with nervous system transmission

DOSE:

Protocol	IO	Repeat?	Notes
Vascular Access (154)	Adult: 40 mg IO , Peds: 0.5 mg/kg IO not to exceed 40 mg	Yes, 20 mg slow IO after IO has initially been flushed with saline. Peds dose 0.25 mg/kg max 20 mg	Infuse slowly over 2 min, then allow to dwell in IO space for 1 min

CONTRAINDICATIONS: Hypersensitivity to the drug

ADVERSE REACTIONS: Seizures; respiratory depression; dizziness; restlessness; confusion; tinnitus; blurred vision; numbness; muscle twitching; hypotension; bradycardia; heart block; nausea; vomiting

LIDOCAINE (Xylocaine) 2% LUBRICANT

CLASS: Topical Anesthetic

ACTION: Produces anesthesia by interfering with nervous system transmission

DOSE:

Protocol	Topical	Notes
Ventilation Management (69)	As Needed	For nasotracheal intubation
Endotracheal Intubation(143)	As Needed	For nasotracheal intubation

CONTRAINDICATIONS: Hypersensitivity to the drug

ADVERSE REACTIONS: Seizures; respiratory depression; dizziness; restlessness; confusion; tinnitus; blurred vision; numbness; muscle twitching; hypotension; bradycardia; heart block; nausea; vomiting

MAGNESIUM SULFATE

CLASS: Electrolyte

ACTION: Membrane stabilization; raises seizure threshold

MAGNESIUM SULFATE (continued)

DOSE:

Protocol	IV/IO	Notes
Obstetrical Emergency (43)	2 - 4 g IV in 50 cc NS over 10 min	2 g dose for pre-eclamptic, 4 g dose for pregnant patient in seizure
Respiratory Distress (51)	2 g IV in 50 cc NS over 10 min	
Tachycardia/Stable (65)	2 g IV in 50 cc NS over 10 min	
Tachycardia/Unstable (67)	2 g IV in 50 cc NS over 10 min	
Peds Tachycardia/Stable (112)	25 mg/kg IV/IO in 50 cc NS over 10 min	Need Physician Orders
Peds Tachycardia/Unstable (114)	25 mg/kg IV/IO in 50 cc NS over 10 min	Need Physician Orders

CONTRAINDICATIONS: Hypersensitivity to the drug; high degree heart block; renal failure

ADVERSE REACTIONS: Hypotension; asystole; respiratory depression; weakness

METOCLOPRAMIDE (Reglan)

CLASS: Antiemetic

ACTION: Dopamine agonist that works by blocking the CNS vomiting chemoreceptor trigger zone (CRT)

DOSE:

Protocol	IV/IO/IM	Notes
Abdominal/Flank Pain, Nausea & Vomiting (15)	10 mg slow IV bolus over 1-2 min OR IM	
Chest Pain(Non Traumatic) (29)	10 mg slow IV bolus over 1-2 min OR IM	
Pain Management (47)	10 mg slow IV bolus over 1-2 min OR IM	
STEMI (61)	10 mg slow IV bolus over 1-2 min OR IM	
Peds Abdominal Pain, Nausea & Vomiting (76)	5 mg slo IV/IO bolus over 1-2 min OR IM	8 years of age and older
Peds Pain Management (102)	5 mg slo IV/IO bolus over 1-2 min OR IM	8 years of age and older

CONTRAINDICATIONS: Known Hypersensitivity

ADVERSE REACTIONS: Restlessness; hyperactivity; anxiety; sedation; increased GI motility; use caution in suspected bowel obstruction. Extra-pyramidal reactions have been seen days to hours after treatment. Protect medication from light (photosensitive)

MIDAZOLAM (Versed)

CLASS: Anxiolytic

ACTION: CNS Depressant

DOSE:

Protocol	IN/IM/IV/IO	Repeat?	Notes
Behavioral Emergency (21) Obstetrical Emergency (43) Seizure (53) Ventilation Management (69) Pediatric Seizure (106) Electrical Therapy/Synchronized Cardioversion (140) Electrical Therapy/Transcutaneous Pacing (141)	0.1 mg/kg IN/IM/IV/IO, max 5mg	Yes, may repeat X 1 after 5 min at 0.05 mg/kg, max 2.5 mg	Call for further dose orders
Peds Ventilation Management (116)	0.1 mg/kg IN/IV/IO, max dose 5 mg	No	Given slowly over 3-5 min, titrate to effect
Endotracheal Intubation (142)	0.1 mg/kg IN/IM/IV/IO, max dose 5 mg	Yes, may repeat X 1 after 5 min at 0.05 mg/kg, max 2.5 mg	For < 12 years of age only, give slowly over 3-5 min, titrate to effect

CONTRAINDICATIONS: Hypersensitivity to the drug; hypotension; clinical signs of shock

ADVERSE REACTIONS: CNS depression; hypotension; respiratory depression

MORPHINE SULFATE

CLASS: Narcotic

ACTION: CNS depressant

DOSE:

Protocol	IM/IV/IO	Repeat?
Pain Management (47)	0.1 mg/kg IM/IV/IO, max single dose 10 mg	Yes, q 10 min until pain is relieved or respiratory
Peds Pain Management (102)	0.1 mg/kg IM/IV/IO, max single dose 10 mg	No
Electrical Therapy/Synchronized Cardioversion (140)	Up to 0.1 mg/kg slow IV/IO, max single dose 10 mg	Yes, q 10 min until pain is relieved or respiratory depression occurs
Electrical Therapy/ Transcutaneous Pacing (141)	Up to 0.1 mg/kg slow IV/IO, max single dose 10 mg	Yes, q 10 min until pain is relieved or respiratory depression occurs

MORPHINE SULFATE (continued)

CONTRAINDICATIONS: Hypersensitivity to opiates; head injuries; chest or abdominal injury; clinical signs of shock

ADVERSE REACTIONS: Respiratory depression; nausea; vomiting; bradycardia; orthostatic hypotension; altered level of consciousness

NALOXONE HYDROCHLORIDE (Narcan)

CLASS: Narcotic antagonist

ACTION: Reverses the effects of narcotics

DOSE:

Protocol	IM/IV/IO	IN	Repeat?
Altered Mental Status/Syncope (20)	0.4 - 2 mg IM/IV/IO, max dose 10 mg	2-4 mg IN	Yes
Overdose/Poisoning (45)	0.4 - 2 mg IM/IV/IO, max dose 10 mg	2-4 mg IN	Yes
Peds Altered Mental Status (81)	0.1 mg/kg IM/IV, max single dose 2 mg, total dose 10 mg	0.1 mg/kg IN, max single dose 2 mg, total dose 10 mg	Yes
Peds Overdose/Poisoning (100)	0.1 mg/kg IM/IV, max single dose 2 mg, total dose 10 mg	2-4 mg IN	Yes

CONTRAINDICATIONS: Patients with a history of hypersensitivity to the drug; intubated patients; the newly born during initial resuscitation

ADVERSE REACTIONS: Rapid administration causes projectile vomiting

NITROGLYCERIN

CLASS: Vasodilator

ACTION: Dilates systemic arteries and veins; reduces both preload and afterload

DOSE:

Protocol	SL	Repeat
Chest Pain (Non Traumatic) (29)	0.4 mg SL	Yes, q 5 min X2
Pulmonary Edema/CHF (49)	0.4 mg SL	Yes, q 5 min as long as HR >60 and systolic BP >100 mm Hg
STEMI (61)	0.4 mg SL	Yes, q 5 min X2

CONTRAINDICATIONS: Hypotension (do not administer if systolic pressure below 100 mm Hg unless ordered by a physician); use of erectile dysfunction medications within the last 48 hours; patients with demonstrated hypersensitivity to nitrates or nitrites

ADVERSE REACTIONS: hypotension

ONDANSETRON HYDROCHLORIDE (Zofran)

CLASS: Selective serotonin blocking agent

ACTION: Antiemetic

DOSE:

Protocol	ODT/IM/IV/IO
Abdominal/Flank Pain, Nausea & Vomiting (15)	4 mg ODT/IM/IV/IO
Chest Pain (Non Traumatic)(29)	4 mg ODT/IM/IV/IO
Pain Management (47)	4 mg ODT/IM/IV/IO
STEMI (61)	4 mg ODT/IM/IV/IO
Peds Abdominal Pain (76)	0.15 mg/kg ODT/IM/IV/IO, up to max dose 4 mg
Peds Pain Management (102)	0.15 mg/kg ODT/IM/IV/IO, up to max dose 4 mg

CONTRAINDICATIONS: Patients with a known hypersensitivity to the drug

ADVERSE REACTIONS: Headache; chest pain; dizziness; hypotension

OXYMETAZOLINE (Afrin) 0.05% SPRAY

CLASS: Sympathomimetic

ACTION: Direct local vasoconstriction

DOSE:

Protocol	IN	Notes
Epistaxis (37)	2 sprays to each nostril	follow with direct pressure
Ventilation Management (69)	1-2 sprays each nostril	for nasotracheal intubation preparation
Peds Epistaxis (94)	2 sprays to each nostril	follow with direct pressure
Endotracheal Intubation (143)	2 sprays to each nostril	for nasotracheal intubation preparation

CONTRAINDICATIONS: Monoamine oxidase inhibitor (MAOI) use within 14 days

ADVERSE REACTIONS: None

PHENYLEPHRINE (Neo-Syneprine)

CLASS: Sympathomimetic

ACTION: Direct local vasoconstriction

DOSE:

Protocol	IN	Notes
Epistaxis (37)	2 sprays to each nostril	follow with direct pressure
Ventilation Management (69)	1-2 sprays each nostril	for nasotracheal intubation preparation
Peds Epistaxis (94)	2 sprays to each nostril	follow with direct pressure
Endotracheal Intubation (143)	1-2 sprays to each nostril	for nasotracheal intubation preparation

CONTRAINDICATIONS: Ventricular tachycardia; severe coronary artery disease; head injured patients with altered mental status

ADVERSE REACTIONS: None

PHENYLEPHRINE PUSH DOSE (injectable)

CLASS: Sympathomimetic

ACTION: Alpha-1 adrenergic receptor agonist

DOSE:

Protocol	IV/IO	Repeat?	Preparation
Bradycardia (23)	100-200 mcg IV/IO	Yes, q 2-5 min to maintain SBP >90	Mix 10 mg phenylephrine into a 100 ml bag of Normal Saline = 100 ml of phenylephrine at a concentration of 100 mcg/ml
Pulmonary Edema/CHF (49)	100-200 mcg IV/IO	Yes, q 2-5 min to maintain SBP >90	Mix 10 mg phenylephrine into a 100 ml bag of Normal Saline = 100 ml of phenylephrine at a concentration of 100 mcg/ml
Sepsis (55)	100-200 mcg IV/IO	Yes, q 2-5 min to maintain SBP >90	Mix 10 mg phenylephrine into a 100 ml bag of Normal Saline = 100 ml of phenylephrine at a concentration of 100 mcg/ml
Shock (57)	100-200 mcg IV/IO	Yes, q 2-5 min to maintain SBP >90	Mix 10 mg phenylephrine into a 100 ml bag of Normal Saline = 100 ml of phenylephrine at a concentration of 100 mcg/ml

CONTRAINDICATIONS: Hypovolemic shock is a relative contraindication. Hypotension due to hypovolemia or distributive shock should be addressed with a fluid bolus before administering Push Dose Phenylephrine.

ADVERSE REACTIONS: None

PROCHLORPERAZINE (Compazine)

CLASS: Antiemetic

ACTION: Dopamine agonist with antiemetic actions

DOSE:

Protocol	IV/IO/IM
Abdominal/Flank Pain, Nausea & Vomiting (15)	Up to 10 mg IV/IM/IO
Chest Pain (Non Traumatic) (29)	Up to 10 mg IV/IM/IO
Pain Management (47)	Up to 10 mg IV/IM/IO
STEMI (61)	Up to 10 mg IV/IM/IO

CONTRAINDICATIONS: Known hypersensitivity

ADVERSE REACTIONS: Possible dystonic reactions

SODIUM BICARBONATE

CLASS: Alkalinizing agent

ACTION: Increases blood pH

DOSE:

Protocol	IV/IO	Repeat?
Hyperkalemia (41)	1.0 mEq/kg slow IV/IO push	No
Overdose/Poisoning (45)	1.0 mEq/kg IV/IO of 8.4% solution	Yes, X1 in 3-5 min
Peds Overdose/Poisoning (100)	1.0 mEq/kg IV/IO of 8.4% solution	Use 4.2% for neonate

CONTRAINDICATIONS: Alkalotic states; respiratory acidosis

ADVERSE REACTIONS: None

SOLU-CORTEF (Hydrocortisone Sodium Succinate)

CLASS: Corticosteroid

ACTION: Anti-inflammatory, replaces absent glucocorticoids, suppresses immune response

DOSE: Assist the patient in administering his or her own Solu-Cortef exactly as prescribed

CONTRAINDICATIONS: Systemic fungal infections, hypersensitivity to the drug

ADVERSE REACTIONS: ECG changes, hypertension, headache

First Response Low-Risk Alpha Evaluate and Release Form (example)

Incident #:

EMS Agency:

Patient Name:

Date of Birth:

Address:

Phone #:

Responding Ambulance Service: **AMR** **MW** **Community Ambulance**

Medical Priority Dispatch System Code:

Time of Patient Contact:

Patient Complaint(s):

Primary Survey Complete

Secondary Survey Complete

Vital Signs: **HR** _____ **RR** _____ **BP** _____ **SPO2** _____ **BG (as applicable)** _____

General Impression:

Confirm the following:

Normal vital signs Time last taken:

Patient does not meet Trauma Field Triage Criteria

No indication for an ECG or cardiac monitoring per SNHD Emergency Medical Care Protocols

Patient is not a threat to self or others

Patient denies AND no evidence or suspicion of the following:

Pregnancy Head trauma Chest pain

Active bleeding Seizure Stroke

Allergic reaction Shortness of breath Syncope, near syncope, or dizziness

Overdose or ingestional error Abdominal pain/flank pain above the
Umbilicus if >35 years of age

Inclusion Criteria:

Call coded and dispatched using MPDS, Alpha or Omega category

Patient between 18 and 65 years of age

Patient has decision making capacity

Patient has a phone to call 9-1-1 if their condition worsens

We have assessed and examined you and have determined your condition as NOT THREATENING TO LIFE/LIMB. With your permission, we will return to service so we may be available in the case of another emergency. An ambulance is continuing to respond. If your condition worsens in any way, call 9-1-1.

I consent to waiting for the ambulance and understand that I can call 9-1-1 if I get worse in any way.

Patient signature

Provider signature

SAMPLE RELEASE OF MEDICAL ASSISTANCE

1. I (or my guardian) have been informed of the reason I should go to the hospital for further emergency care.
2. I (or my guardian) have been informed that only an initial evaluation has been rendered to me and have been advised that I seek the advice of a physician as soon as possible.
3. I (or my guardian) have been informed of the potential consequences and/or complications that may result in my (or my guardian's) refusal to go to the hospital for further emergency care.
4. I (or my guardian), the undersigned, have been advised that emergency medical care on my/the patient's behalf is necessary, and that refusal of recommended care and transport to a hospital facility may result in death, or imperil my/the patient's health by increasing the opportunity for consequences or complications. Nevertheless, and understanding all of the above, I (or my guardian), refuse to:


- accept emergency medical care
- accept transport to a hospital facility
- accept transport to _____ Hospital as directed by Southern Nevada EMS protocols, but request transport to _____ Hospital; and

assume all risks and consequences resulting from my (or my guardian's) decision, and release Clark County provider agencies, and all personnel directly or indirectly involved in my care from any and all liability resulting from my (or my guardian's) refusal. I have had the opportunity to ask all of the questions I feel necessary to provide this informed refusal.

5. The reason for this refusal is as follows: (to be completed by patient/guardian) _____

Patient's Name:	DOB:	
Patient's Address:		
Patient's Phone Number:		
Signature (Patient/Guardian):		
Witness:		
Witness:		
Date:	Time:	Incident #:
Refused to Sign (Patient/Guardian):		
Telemetry Physician:	Hospital:	

Sample Algorithm, Release of Medical Assistance



Note:

1. For all patients refusing transport who meet Trauma Field Triage Criteria protocol, contact a trauma center.
2. EMS personnel may make telemetry contact for further guidance at any time.

Scope of Practice

Southern Nevada Health District
Office of Emergency Medical Services & Trauma System
Authorized Medication List

The following is the formulary used by EMS agencies in Clark County. Licensed EMS providers working in the EMS System for a permitted agency are authorized, within their level of certification and training, to administer medications as directed by the written treatment protocols.

Medications	EMT	AEMT	Paramedic
Acetylsalicylic Acid	X	X	X
Adenosine			X
Albuterol		X	X
Amiodarone			X
Atropine Sulfate			X
Bronchodilator Metered Dose Inhaler	X	X	X
Calcium Chloride			X
Diazepam			X
Diphenhydramine Hydrochloride		X	X
Dopamine Hydrochloride			X
Droperidol			X
Epinephrine		X	X
Epinephrine Autoinjector	X	X	X
Etomidate			X
Fentanyl Citrate			X
Glucagon		X	X
Glucose - Oral	X	X	X
Glucose Sterile Injectible		X	X
Hydromorphone			X
Hydroxocobalamin			X
Ipratropium Bromide			X
Ipratropium Bromide & Albuterol Sulfate			X
Ketamine			X
Lidocaine			X
Magnesium Sulfate			X
Metoclopramide			X
Midazolam			X
Morphine Sulfate			X
Naloxone Hydrochloride	X	X	X
Nitroglycerin	X	X	X
Ondansetron Hydrochloride		X	X
Oxymetazoline	X	X	X
Phenylephrine	X	X	X
Push Dose Phenylephrine			X
Prochlorperzine			X
Sodium Bicarbonate			X
Solu-Cortef		X	X

Scope of Practice

**Southern Nevada Health District
Office of Emergency Medical Services & Trauma System
Authorized Skills List**

The following are the authorized skills used by EMS providers in Clark County. Licensed EMS providers working in the EMS System for a permitted agency are authorized, within their level of certification and training, to perform the skills as directed by the written treatment protocols.

Skills	EMT	AEMT	Paramedic
12-Lead ECG Interpretation			X
3-Lead ECG Interpretation			X
Airway Adjunct OPA/NPA	X	X	X
Airway Suction	X	X	X
Capnometry (Color Change Device)		X	X
Capnometry (Continuous Waveform)			X
Carotid Massage			X
Cervical Stabilization	X	X	X
CPAP		X	X
CPR	X	X	X
Defibrillation - AED	X	X	X
Defibrillation - Manual			X
Endotracheal Intubation - Nasal			X
Endotracheal Intubation - Oral			X
Extraglottic Airway		X	X
Gastric Decompression			X
Glucose Measurement	X	X	X
Hemorrhage Control	X	X	X
Medication Administration	X	X	X
Needle Cricothyroidotomy			X
Needle Thoracentesis			X
Oxygen Administration	X	X	X
Patient Assessment	X	X	X
Pulse Oximetry	X	X	X
Restraints	X	X	X
Splinting	X	X	X
Stroke Screening	X	X	X
Synchronized Cardioversion			X
Thermometer	X	X	X
Tracheostomy Tube Replacement			X
Transcutaneous Pacing			X
Vagal Maneuvers			X
Vascular Access IV		X	X
Vascular Access IO		X (Adult)	X

Southern Nevada Health District
Office of Emergency Medical Services and Trauma System
SUBJECT: Receiving Hospital Directory

This directory is subject to change and is provided for informational purposes only

HOSPITAL NAME ADDRESS PHONE	HOSP CODE	TRAUMA LEVEL	BURN CENTER	STROKE DEST	PEDS DEST	L & D	HELIP AD	SPECIAL SERVICE S
Boulder City Hospital 901 Adams Blvd Boulder City, NV 89005 (702) 294-5751	H-8						X	
Centennial Hills Hospital 6900 North Durango Dr Las Vegas, NV 89149-4409 (702) 359-9700	H-19			X		X	X	
Desert Springs Hospital 2075 E. Flamingo Rd. Las Vegas, NV 89119 (702) 369-7772	H-2			X			X	
Henderson Hospital 1050 Galleria Dr. Henderson, NV 89011 (702) 963-7000	H-18					X	x	
Mesa View Hospital 1299 Bertha Howe Avenue Mesquite, NV 89027 (702) 346-8040	H-75					X	X	
Mike O'Callaghan Federal Hospital 4700 Las Vegas Blvd N Nellis AFB, NV 89191-6600 (702) 653-2344	H-3	Level III				X		
Mountain View Hospital 3100 N Tenaya Way Las Vegas, NV 89128 (702) 255-5025	H-11			X		X		
North Vista Hospital 1409 East Lake Mead Blvd North Las Vegas, NV 89030 (702) 657-5512	H-5							
Southern Hills Hospital 9300 W. Sunset Road Las Vegas, NV 89148 (702) 880-2800	H-16			X		X	X	
Spring Valley Hospital 5400 South Rainbow Blvd Las Vegas, NV 89118 (702) 853-3611	H-15			X		X	X	
St. Rose San Martin 8280 West Warm Springs Rd Las Vegas, NV 89113 (702) 432-8600	H-73			X		X	X	
St. Rose Siena 3001 St. Rose Pkwy Henderson, NV 89052 (702) 616-5600	H-72 (-T/-P)	Level III Pediatric Level IV		X	X	X	X	
Summerlin Medical Center 657 Town Center Dr. Las Vegas, NV 89144 (702) 233-7033	H-1 (-P)			X	X	X	X	
Sunrise Hospital 3186 South Maryland Pkwy Las Vegas, NV 89109 (702) 731-8585	H-6 (-T/-P)	Level II Pediatric Level III and IV	X	X	X	X	X	SA Evaluati on (Patient
University Medical Center 1800 W. Charleston Blvd Las Vegas, NV 89102 (702) 383-2000M, 3969Trauma	H-4 (-T/-P)	Level I Pediatric Level II	X	X	X	X	X	SA Evaluati on (Patient
Valley Hospital 620 Shadow Lane Las Vegas,, NV 89106 (702) 388-4506	H-9			X			X	

Southern Nevada Health District									
Office of Emergency Medical Services and Trauma System									
SUBJECT: Receiving Hospital Directory									
This directory is subject to change and is provided for									
informational purposes only									
HOSPITAL NAME ADDRESS PHONE	Remote Outpatient ED Alternate Destination Protocol	Public Intoxication Protocol							
Crossroads of Southern Nevada 2121 W. Charleston (702)382-7746		X							
Dignity Blue Diamond 4855 Blue Diamond (702)407- 0725	X								
Dignity North Las Vegas 1550 W. Craig (702)642-0760	X								
Dignity Sahara 4980 W. Sahara (702)257- 0824	X								
Dignity West Flamingo 9880 W. Flamingo (702)243- 1795	X								
Elite Medical 150 E Harmon Ave (702)546-0911	X								
ER at Alliante 7207 N. Aliante Pkwy (702)962-5100	X								
ER at Blue Diamond 9217 S Cimarron Rd (702) 853-3611	X								
ER at Green Valley Ranch 2581 St Rose Parkway (702) 780-2700	X								
ER at the Lakes 3325 S. Fort Apache (702)962-0500	X								
ER at Skye Canyon 9860 Skye Canyon Park Dr (702)962-0700	X								
ER at South Las Vegas Blvd 10770 South Las Vegas Blvd (702)962-0800	X								
ER at Valley Vista 7230 N Decatur Blvd (702)534-7200	X								
St Rose De Lima 102 E Lake Mead Pkwy Henderson, NV 89015 (702)616-460	X								
Westcare 323 North Maryland Pkwy 702-385-3300		X							

Telemetry Radio Map

CCFD-2019 FLEETMAP				ALL OTHER FIRE DEPTS (LVFD, NLVFD, HFD, BCFD)				EMS (AMR, MWA, CA, GEMS, etc...)			
Zone #	Zone 12	Zone 13	Zone 14	Zone #	Zone 12	Zone 13	Zone 14	Zone #	ZONE-5	ZONE-6	ZONE-7
Zone Name	Hospitals 1	Hospitals 2	Hospitals 3	Zone Name	Hospitals 1	Hospitals 2	Hospitals 3	Zone Name	MEDICAL_ZONE	HOSPITAL_1	HOSPITAL_2
Chn	Talk Group	Talk Group	Talk Group	Chn	Talk Group	Talk Group	Talk Group	Chn	Talk Group	Talk Group	Talk Group
1	ALIANTE	SR SIENA	NEW HOSP 6	1	ER at HCA	SR SIENA	NEW HOSP 6	1	AMR	BOULDER_CITY	SUMMERLIN
2	BLDR CITY	SUMRLIN	NEW HOSP 7	2	BLDR CITY	SUMRLIN	NEW HOSP 7	2	MEDIC_WEST	CENTENNIAL	SUMRLIN_PEDS
3	CENTEN HILLS	SUMRLIN PED	NEW HOSP 8	3	CENTEN HILLS	SUMRLIN PED	NEW HOSP 8	3	COMMUNITY	DSRT_SPRINGS	SUNRISE_ER
4	CROSSROADS	SUN ER	NEW HOSP 9	4	DETOX LOC	SUN ER	NEW HOSP 9	4	MED_BRANCH	MESA_VIEW	SUNRISE_PEDS
5	DSRT SPRG	SUN PEDS	NEW HOSP 10	5	DSRT SPRG	SUN PEDS	NEW HOSP 10	5	ALL_HOSPITLS	MTN_VIEW	SUNRISE_TRAUMA
6	GREEN VALLEY	SUN TRMA	NEW HOSP 11	6	ER at VHS	SUN TRMA	NEW HOSP 11	6	EMS_TAC_6	NELLIS/CALAGHN	UMC_ER
7	HENDERSON	UMC ER	NEW HOSP 12	7	HENDERSON	UMC ER	NEW HOSP 12	7	EMS_TAC_7	NORTH_VISTA	UMC_PEDS
8	LAKES ER	UMC PEDS	NEW HOSP 13	8	BLANK	UMC PEDS	NEW HOSP 13	8	EMS_TAC_8	SOUTH_HILLS	UMC_TRAUMA
9	MESA VIEW	UMC TRMA	NEW HOSP 14	9	MESA VIEW	UMC TRMA	NEW HOSP 14	9	EMS_TAC_9	SPRG_VALLEY	VALLEY
10	MTN VIEW	VALLEY	NEW HOSP 15	10	MTN VIEW	VALLEY	NEW HOSP 15	10	EMS_TAC_10	ST_ROSE_LIMA	VA_MC
11	NELLIS	VAMC	NEW HOSP 16	11	NELLIS	VAMC	NEW HOSP 16	11	EMS_TAC_11	SAN_MARTIN	NEW_HOSP_1
12	N VISTA	NEW HOSP 1	NEW HOSP 17	12	N VISTA	SR MICRO	NEW HOSP 17	12	EMS_TAC_12	ST_ROSE_SIEN	NEW_HOSP_2
13	SO. HILLS	NEW HOSP 2	NEW HOSP 18	13	SO. HILLS	NEW HOSP 2	NEW HOSP 18	13	EMS_TAC_13	UHS_HENDERSON	NEW_HOSP_3
14	SPRG VALL	NEW HOSP 3	NEW HOSP 19	14	SPRG VALL	NEW HOSP 3	NEW HOSP 19	14	EMS_TAC_14	LAKES_ER	MERCY_AIR
15	SR DELIMA	NEW HOSP 4	NEW HOSP 20	15	SR DELIMA	NEW HOSP 4	NEW HOSP 20	15	EMS_TAC_15	BLANK	BLANK
16	SR SAN MART	NEW HOSP 5	NEW HOSP 21	16	SR SAN MART	NEW HOSP 5	NEW HOSP 21	16	BLANK	BLANK	BLANK

LEGEND	
SR MICRO	ER at HCA
St. Rose Blue Diamond	aliante ER
St. Rose NLV	the lakes ER
St. Rose Sahara	
St. Rose W. Flamingo	
ELITE MED FED	
ER at VHS	
green valley ER	
blue diamond ER	
DETOX LOC	
crossroads	
westcare	

11/21/2019	Communications to SR MICRO can be performed via zone - 13 (Hospital 2) on channel - 12 (New HOSP 1) via the CCFD FLEETMAP. Communications to SR MICRO can be performed via zone - 13 (Hospital 2) on channel - 12 (SR MICRO) via the ALL OTHER FIRE DEPTS FLEETMAP. Communications to SR MICRO are NOT available at this time via the EMS FLEETMAP.
12/10/2019	Communications to Elite Medical FED can be performed via zone - 13 (Hospital 2) on channel - 13 (New HOSP 2) via the CCFD FLEETMAP. Communications to Elite Medical FED can be performed via zone - 13 (Hospital 2) on channel - 13 (New HOSP 2) via the ALL OTHER FIRE DEPTS FLEETMAP. Communications to Elite Medical FED are NOT available at this time via the EMS FLEETMAP.
12/10/2019	Communications to ER at VHS can be performed via zone - 12 (Hospital 1) on channel - 6 (Green Valley) via the CCFD FLEETMAP. Communications to ER at VHS can be performed via zone - 12 (Hospital 1) on channel - 6 (ER at VHS) via the ALL OTHER FIRE DEPTS FLEETMAP. Communications to ER at VHS can be performed via zone - varies (Hospital 2) on channel - 12 (NEW HOSP 2) via the EMS FLEETMAP.
12/10/2019	Communications to DETOX LOC can be performed via zone - 12 (Hospital 1) on channel - 4 (CROSS ROADS) via the CCFD FLEETMAP. Communications to DETOX LOC can be performed via zone - 12 (Hospital 1) on channel - 4 (DETOX LOC) via the ALL OTHER FIRE DEPTS FLEETMAP. Communications to DETOX LOC are NOT available at this time via the EMS FLEETMAP.
12/10/2019	Communications to ER at HCA can be performed via zone - 12 (Hospital 1) on channel - 1 (ALIANTE) via the CCFD FLEETMAP. Communications to ER at HCA can be performed via zone - 12 (Hospital 1) on channel - 1 (ER at HCA) via the CCFD FLEETMAP. Communications to ER at HCA can be performed via zone - varies (Hospital 2) on channel - 11 (NEW HOSP 1) via the EMS FLEETMAP.

Appendix E

Mass Casualty Incident

Licensed EMS providers who are on duty for a permitted agency may operate within their scope of practice at a receiving facility during a Mass Casualty Incident if that incident is classified as a Level 3, Level 2, Level 1, Extreme MCI or Hostile MCI as defined by the Fire Alarm Office Standard Operating Procedure, M-1 dated 5/09/2016 if requested by an authorized agent of the receiving facility. Further, if the licensed EMS provider is requested through the Incident Commander or Designee to assist with Mass Casualty Incident (as defined above) related patient surge at a hospital to provide triage expertise or act as a liaison with the EMS system they may do so with consideration being made to EMS system needs, status and relevant law.

Revised and approved by MAB 12-05-2018