

| Disease | 2018 | | 2019 | | 2020 | | Rate (Cases per 100,000 per quarter) | | Quarter Rate Comparison |
|---|-------|------|-------|------|-------|------|--------------------------------------|--------------|-----------------------------------|
| | Qtr 1 | YTD | Qtr 1 | YTD | Qtr 1 | YTD | Qtr 1 (2015-2019 aggregated) | Qtr 1 (2020) | Change b/t current & past 5-year? |
| VACCINE PREVENTABLE | | | | | | | | | |
| Haemophilus influenzae, invasive | 5 | 5 | 9 | 9 | 8 | 8 | 0.35 | . | ↑ |
| Hepatitis A | 7 | 7 | 19 | 19 | 9 | 9 | 0.28 | . | ↑ |
| Hepatitis B, acute | 5 | 5 | 3 | 3 | 0 | 0 | 0.16 | . | ↓ |
| Hepatitis B, chronic | 90 | 90 | 101 | 101 | 129 | 129 | 2.13 | 5.69 | ↑X |
| Influenza | 744 | 744 | 683 | 683 | 790 | 790 | 24.76 | 34.83 | ↑X |
| Influenza-associated pediatric mortality | 0 | 0 | 2 | 2 | 0 | 0 | . | . | ↓ |
| Meningococcal disease (<i>N. meningitidis</i>) | 2 | 2 | 1 | 1 | 1 | 1 | . | . | ↓ |
| Mumps | 3 | 3 | 0 | 0 | 0 | 0 | . | . | ↓ |
| Pertussis | 12 | 12 | 18 | 18 | 5 | 5 | 0.73 | . | ↓ |
| SEXUALLY TRANSMITTED | | | | | | | | | |
| Chlamydia | 3295 | 3295 | 3422 | 3422 | 3351 | 3351 | 137.20 | 147.75 | ↑X |
| Gonorrhea | 1241 | 1241 | 1270 | 1270 | 1189 | 1189 | 45.63 | 52.43 | ↑X |
| HIV | 105 | 105 | 110 | 110 | 53 | 53 | 4.58 | . | ↓ |
| Stage 3 HIV (AIDS) | 30 | 30 | 29 | 29 | 26 | 26 | 1.70 | . | ↓ |
| Syphilis (Primary & Secondary) | 145 | 145 | 147 | 147 | 167 | 167 | 4.90 | 7.36 | ↑X |
| Syphilis (Early non-primary, non-secondary) | 98 | 98 | 108 | 108 | 90 | 90 | 4.96 | 3.97 | ↓ |
| CONGENITAL CONDITIONS | | | | | | | | | |
| Congenital Syphilis | 3 | 3 | 4 | 4 | 12 | 12 | 0.16 | 0.53 | ↑X |
| Hepatitis C, Perinatal Infection | 0 | 0 | 1 | 1 | 0 | 0 | . | . | ↓ |
| ENTERICS | | | | | | | | | |
| Amebiasis | 1 | 1 | 4 | 4 | 3 | 3 | . | . | ↑ |
| Campylobacteriosis | 19 | 19 | 31 | 31 | 23 | 23 | 1.13 | 1.01 | ↓ |
| Cryptosporidiosis | 0 | 0 | 1 | 1 | 4 | 4 | . | . | ↑ |
| Giardiasis | 14 | 14 | 10 | 10 | 8 | 8 | 0.47 | . | ↓ |
| Rotavirus | 3 | 3 | 6 | 6 | 12 | 12 | 0.70 | 0.53 | ↓ |
| Salmonellosis | 25 | 25 | 18 | 18 | 23 | 23 | 1.14 | 1.01 | ↓ |
| Shiga toxin-producing <i>E. coli</i> (STEC) | 2 | 2 | 4 | 4 | 8 | 8 | 0.27 | . | ↑ |
| Shigellosis | 20 | 20 | 11 | 11 | 10 | 10 | 0.54 | . | ↓ |
| Vibriosis (Non-cholera <i>Vibrio</i> species infection) | 0 | 0 | 1 | 1 | 2 | 2 | . | . | ↑ |
| Yersiniosis | 0 | 0 | 1 | 1 | 1 | 1 | . | . | ↑ |
| OTHER | | | | | | | | | |
| Coccidioidomycosis | 63 | 63 | 20 | 20 | 33 | 33 | 1.35 | 1.46 | ↑ |
| Encephalitis | 1 | 1 | 0 | 0 | 0 | 0 | . | . | ↓ |
| Exposure, Chemical or Biological | 3 | 3 | 5 | 5 | 0 | 0 | . | . | ↓ |
| Hepatitis C, acute | 2 | 2 | 4 | 4 | 0 | 0 | 0.16 | . | ↓ |
| Hepatitis C, chronic | 626 | 626 | 1708 | 1708 | 1335 | 1335 | 21.38 | 58.86 | ↑X |
| Invasive Pneumococcal Disease | 89 | 89 | 83 | 83 | 101 | 101 | 3.16 | 4.45 | ↑X |
| Lead Poisoning | 48 | 48 | 32 | 32 | 40 | 40 | 1.42 | 1.76 | ↑ |
| Legionellosis | 4 | 4 | 3 | 3 | 6 | 6 | 0.18 | . | ↑ |
| Listeriosis | 1 | 1 | 3 | 3 | 0 | 0 | . | . | ↓ |
| Lyme Disease | 5 | 5 | 3 | 3 | 2 | 2 | 0.13 | . | ↓ |
| Malaria | 0 | 0 | 1 | 1 | 0 | 0 | . | . | ↓ |
| Meningitis, Aseptic | 9 | 9 | 8 | 8 | 25 | 25 | 0.32 | 1.10 | ↑X |
| Meningitis, Bacterial Other | 9 | 9 | 7 | 7 | 7 | 7 | 0.33 | . | ↓ |
| Meningitis, Fungal | 3 | 3 | 3 | 3 | 1 | 1 | . | . | ↓ |
| RSV | 1200 | 1200 | 1698 | 1698 | 1256 | 1256 | 49.60 | 55.38 | ↑X |
| Spotted Fever Rickettsiosis | 1 | 1 | 0 | 0 | 0 | 0 | . | . | ↓ |
| Streptococcal Toxic Shock Syndrome (STSS) | 13 | 13 | 11 | 11 | 10 | 10 | 0.34 | . | ↑ |
| Tuberculosis, Active | 12 | 12 | 7 | 7 | 10 | 10 | 0.41 | . | ↑ |

*Use of illness in data aggregation for cases other than STD or TB (since Jan-2013) causes changes in cases reported here from previously released reports. Numbers are provisional including confirmed, probable, and suspect cases that are reportable to CDC. HIV/AIDS/TB case counts are provided on a quarterly basis. Rate suppression denoted by '.' for rates corresponding to case counts < 12.

~Diseases not reported in the past five years (aggregate data) and not reported during the current reporting period are not included in this report.

--Confidence intervals (not shown) for the quarterly disease incidence rates provided a basis for an informal statistical test to determine if the current quarterly rates changed significantly from those of the previous 5-year aggregated rates. Green text represents rates that decreased significantly, whereas red text represents rates that increased significantly. Statistically significant changes are indicated by 'X.'